

Individuals using assistive technology may not be able to fully access the information contained in this file. For assistance, please send an e-mail to: ocod@fda.hhs.gov and include 508 Accommodation and the title of the document in the subject line of your e-mail.

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use CUTAQUIG safely and effectively. See full prescribing information for CUTAQUIG.

CUTAQUIG® (Immune Globulin Subcutaneous (Human) - hipp), 16.5% solution
Initial U.S. Approval: 2018

<p>WARNING: THROMBOSIS</p> <p><i>See full prescribing information for complete boxed warning</i></p> <ul style="list-style-type: none"> Thrombosis may occur with immune globulin products, including CUTAQUIG. Risk factors may include: advanced age, prolonged immobilization, hypercoagulable conditions, history of venous or arterial thrombosis, use of estrogens, indwelling vascular catheters, hyperviscosity, and cardiovascular risk factors. For patients at risk of thrombosis, administer CUTAQUIG at the minimum dose and infusion rate practicable. Ensure adequate hydration in patients before administration. Monitor for signs and symptoms of thrombosis and assess blood viscosity in patients at risk of hyperviscosity.

Infusion Parameters*		Infusion Number	
		Initial 2 infusions	Subsequent infusions
Rate (mL/hr/site)			weeks up to a maximum of 29
	2-6 years	≤ 10	gradually increase by approximately 5-10 mL/site every 2-4 weeks up to a maximum of 15.5
	Adults ≥17 years	≤ 20	gradually increase by approximately 10 mL/hr/site every 2-4 weeks up to a maximum of 52
	Children 2-17 years	≤ 15	gradually increase by approximately 10 mL/hr/site every 2-4 weeks up to a maximum of 25

* As tolerated

-----RECENT MAJOR CHANGES-----

Indications and Usage (1)	10/2021
Dosage and Administration, Dose (2.1)	10/2021
Dosage and Administration, Administration (2.3)	10/2021

-----INDICATIONS AND USAGE-----

CUTAQUIG (Immune Globulin Subcutaneous (Human) - hipp) is a 16.5% immune globulin solution for subcutaneous infusion indicated for treatment of primary humoral immunodeficiency (PI) in adults and pediatric patients 2 years of age and older. (1)

-----DOSAGE AND ADMINISTRATION-----

For subcutaneous use only

Before switching to CUTAQUIG, obtain the patient's serum IgG trough level to guide subsequent dose adjustments.

Dose (2.1)

- Switching from Immune Globulin Intravenous (IGIV) to CUTAQUIG: calculate the dosing by using a dose conversion factor (1.30); Switching from other Immune Globulin Subcutaneous (IGSC): dosing should be the same as for previous IGSC.
- Weekly: Start CUTAQUIG one week after last IGIV infusion

$$\text{Initial weekly dose} = \frac{\text{Previous IGIV dose (in grams)} \times 1.30}{\text{No. of weeks between IGIV doses}}$$

- Every other week: Start CUTAQUIG 1 or 2 weeks after the last IGIV dosing or 1 week after the last IGSC administration. Infuse twice the calculated weekly dose.
- More frequent than weekly (2-7 times a week): start CUTAQUIG 1 week after last IGIV or IGSC administration. Divide the calculated weekly dose by the desired number of infusions per week.
- Adjust dosing according to patient's pharmacokinetics and clinical response. (see [Dose Adjustment](#))

Administration (2.3):

Administer at regular interval from daily up to every other week. Infusion rate and infusion volume per site may gradually be increased every 2-4 weeks as tolerated by patients.

Infusion Parameters*		Infusion Number	
		Initial 2 infusions	Subsequent infusions
Volume (mL/site)	Adults ≥17 years	≤ 25	gradually increase by approximately 10 mL/site every 2-4 weeks up to a maximum of 40
	7-17 years	≤ 15	gradually increase by approximately 5-10 mL/site every 2-4

Infusion sites: up to 6 infusion sites simultaneously, with at least 2 inches between sites. Rotate sites for each administration. (2.3)

-----**DOSAGE FORMS AND STRENGTHS**-----

CUTAQUIG is a solution containing 16.5% IgG (165 mg/mL) for subcutaneous infusion. (3)

-----**CONTRAINDICATIONS**-----

- History of anaphylactic or severe systemic reaction to human immune globulin or other components of CUTAQUIG (Polysorbate 80). (4)
- IgA deficient patients with antibodies against IgA and a history of hypersensitivity. (4)

-----**WARNINGS AND PRECAUTIONS**-----

- Hypersensitivity and anaphylactic reactions can occur with CUTAQUIG. IgA-deficient patients with anti-IgA antibodies are at greater risk of severe reactions. (5.1)
- Thrombosis may occur with CUTAQUIG. (5.2)
- Falsely elevated blood glucose readings may occur during and after the infusion of CUTAQUIG with some glucometer and test strip systems. (5.3)
- Aseptic meningitis syndrome (AMS) may occur within 2 days of treatment. (5.4)
- Monitor patients for signs and symptoms of renal dysfunction. Monitor blood urea nitrogen, serum creatinine, and urine output in patients at risk of acute renal failure. (5.5)

- Monitor for clinical signs and symptoms of hemolysis, especially in patients with pre-existing anemia and/or cardiovascular or pulmonary compromise. (5.6)
- Monitor for pulmonary adverse reactions (transfusion-related acute lung injury, TRALI). (5.7)
- CUTAQUIG is made from human plasma and may carry a risk of transmitting infectious agents, e.g. viruses, the variant Creutzfeldt-Jakob disease (vCJD) agent, and, theoretically, the Creutzfeldt-Jakob disease (CJD) agent. (5.8)

-----**ADVERSE REACTIONS**-----

The most common adverse reactions (≥ 5% of study subjects) were local infusion site reactions (such as redness, swelling, itching), headache, fever, dermatitis, asthma, diarrhea, and cough. (6)

To report SUSPECTED ADVERSE REACTIONS, contact Pfizer Inc. at 1-800-438-1985 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

-----**DRUG INTERACTIONS**-----

The passive transfer of antibodies may transiently interfere with the immune response to live virus vaccines, such as measles, mumps, varicella and rubella (7.2) and may lead to misinterpretation of the results of serological testing. (7.1)

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Revised: 10/2021

FULL PRESCRIBING INFORMATION: CONTENTS*

WARNING: THROMBOSIS

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

- 2.1 Dose
- 2.2 Preparation
- 2.3 Administration

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Hypersensitivity
- 5.2 Thrombosis
- 5.3 Falsely Elevated Glucose Readings
- 5.4 Aseptic Meningitis Syndrome (AMS)
- 5.5 Renal Dysfunction/Failure
- 5.6 Hemolysis
- 5.7 Transfusion-related Acute Lung Injury (TRALI)
- 5.8 Transmittable Infectious Agents
- 5.9 Interference with Laboratory Tests

6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience
- 6.2 Postmarketing Experience

7 DRUG INTERACTIONS

- 7.1 Serological Testing
- 7.2 Live Virus Vaccines

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Lactation
- 8.4 Pediatric Use
- 8.5 Geriatric Use

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
- 13.2 Animal Toxicology and/or Pharmacology

14 CLINICAL STUDIES

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

PATIENT PACKAGE INSERT

*Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

CUTAQUIG (Immune Globulin Subcutaneous (Human) - hipp) 16.5% solution

WARNING: THROMBOSIS

- Thrombosis may occur with immune globulin products, including CUTAQUIG. Risk factors may include: advanced age, prolonged immobilization, hypercoagulable conditions, history of venous or arterial thrombosis, use of estrogens, indwelling central vascular catheters, hyperviscosity, and cardiovascular risk factors. Thrombosis may occur in the absence of known risk factors. [see [Warnings and Precautions \(5.2\)](#), [Patient Counseling Information \(17\)](#)]
- For patients at risk of thrombosis, administer CUTAQUIG at the minimum dose and infusion rate practicable. Ensure adequate hydration in patients before administration. Monitor for signs and symptoms of thrombosis and assess blood viscosity in patients at risk of hyperviscosity. [see [Warnings and Precautions \(5.2\)](#)]

1 INDICATIONS AND USAGE

CUTAQUIG (Immune Globulin Subcutaneous (Human) - hipp) is a 16.5% immune globulin solution for subcutaneous infusion (IGSC), indicated as replacement therapy for primary humoral immunodeficiency (PI) in adults and pediatric patients 2 years of age and older. This includes, but is not limited to, common variable immunodeficiency (CVID), X-linked agammaglobulinemia, congenital agammaglobulinemia, Wiskott-Aldrich syndrome, and severe combined immunodeficiencies.

2 DOSAGE AND ADMINISTRATION

For subcutaneous use only.

- Before receiving treatment with CUTAQUIG: Obtain the patient's serum Immunoglobulin G (IgG) trough level to guide subsequent dose adjustments (see [Dose Adjustment](#)).

2.1 Dose

- Individualize the dose based on the patient's pharmacokinetic and clinical response. Monitor serum IgG trough levels regularly to guide subsequent dose adjustments as needed (see [Dose Adjustment](#)).
- Start CUTAQUIG treatment one week after the last Immune Globulin Intravenous (IGIV)/ Immune Globulin Subcutaneous (IGSC) infusion.

Dose for patients switching to CUTAQUIG from IGIV:

- Establish the initial weekly dose of CUTAQUIG by converting the monthly IGIV dose into an equivalent weekly dose and increasing it using a dose adjustment factor.
 - To calculate the initial weekly dose of CUTAQUIG, divide the monthly IGIV dose in grams by the number of weeks between IGIV infusions and then multiply this value with a Dose Adjustment Factor of 1.30.
$$\text{Initial weekly dose} = \frac{\text{Previous IGIV dose (in grams)} \times 1.30}{\text{Number of weeks between IGIV doses}}$$
 - To convert the dose (in grams) to milliliters (mL), multiply the calculated dose (in grams) by 6.
- Provided the total weekly dose is maintained, any dosing interval from daily up to weekly can be used and will result in systemic IgG exposure that is comparable to the previous IGIV treatment.
- For every other week dosing, multiply the calculated weekly CUTAQUIG dose by 2.
- For more frequent than weekly dosing (2-7 times a week) divide the calculated weekly CUTAQUIG dose by the desired number of infusions per week (e.g.: if you want to infuse CUTAQUIG 2 times per week, divide the weekly dose by 2).

- On average, serum IgG trough levels were approximately 23% higher during CUTAQUIG administration compared to those obtained during prior IGIV therapy.
- To guide dose adjustments, see [Table 1](#) under [Dose Adjustment](#).

Dose for patients switching to CUTAQUIG from IGSC:

- It is recommended to maintain the same weekly dosing (in grams) of CUTAQUIG that was used for the previous Immune Globulin Subcutaneous (Human) (IGSC) therapy (in grams).
- For every other week dosing, multiply the weekly dose by 2.
- For more frequent than weekly dosing (2-7 times a week) divide the weekly dose by the desired number of infusions per week (e.g.: if you want to infusion CUTAQUIG 2 times per week, divide the weekly dose by 2).
- To convert the dose (in grams) to milliliters (mL), multiply the calculated dose (in grams) by 6.
- Obtain a trough IgG level before switching, monitor clinical response and check the trough IgG level 2 to 3 months after initiating CUTAQUIG.
- To guide dose adjustments, see [Table 1](#) under [Dose Adjustment](#), if the trough IgG level during CUTAQUIG administration differs from the trough IgG level obtained during treatment with the previously administered IGSC product or target trough level.

Dose Adjustment:

Over time the dose may need to be individualized for each patient, depending on the pharmacokinetic and clinical response and on the desired IgG trough level. Measure the patient’s serum IgG trough level 2-3 months after switching to CUTAQUIG or after the last CUTAQUIG dose adjustment, in order to determine if a dose adjustment is necessary.

Calculate the difference between the patient’s target serum IgG trough level (in mg/dL) and the IgG trough level obtained during subcutaneous treatment with CUTAQUIG. Find this difference in column 1 of [Table 1](#) and according to frequency of administration (weekly or every other week) locate the corresponding adjustment amount of CUTAQUIG in mL/administration according to the body weight of the patient.

For more frequent dosing than weekly or every other week, add the weekly increment from [Table 1](#) to the weekly equivalent dose of the patient and then divide by the desired number of administrations per week.

Use the patient’s clinical response as the primary point to consider for any dose adjustment. Additional dose increments may be indicated based on the patient’s clinical response (i.e., infection frequency and severity).

Table 1 Incremental adjustment (mL) of weekly or every other week CUTAQUIG dosing based on the calculated difference between actual IgG trough level and the target trough level for the patient*

Difference from Target Serum IgG Trough Level	Dosing Frequency	Weight-adjusted Dose Increment (mL [†])				
		Body weight				
		30 kg	50 kg	70 kg	90 kg	110 kg
50 mg/dl	Weekly	4	6	8	11	13
	Every other week	7	12	16	21	26
100 mg/dl	Weekly	7	12	16	21	26
	Every other week	14	24	33	42	52
200 mg/dl	Weekly	14	24	33	42	52
	Every other week	28	47	66	85	104
300 mg/dl	Weekly	21	35	49	64	78
	Every other week	42	71	99	127	155

*Derived from a linear regression model of trough levels and weekly dose per kg body weight.

For example: a patient with body weight of 70 kg is treated weekly and has a trough level of 600 mg/dL, but the target trough level is 900 mg/dL. The difference between the actual trough level (600) and the targeted trough level (900) is plus 300 mg/dL. Therefore, the recommended increase in the weekly dose would be approximately 49 mL.

A patient with a body weight of 50 kg, on an every other week dosing schedule and with an actual trough level of 900 mg/dL has a target trough level of 700 mg/dL. The difference between actual trough level (900) and targeted trough level (700) is minus 200 mg/dL, therefore the patient needs a decrease in his every other week dose of approximately 47 mL.

Monitor the patient's clinical response and periodically check trough IgG levels, and repeat dose adjustment as needed.

Measles Exposure

If a patient has been exposed to measles, it may be prudent to administer a dose of Immune Globulin Intravenous as soon as possible and within 6 days of exposure. A dose of 400 mg/kg should provide a serum level > 240 mIU/mL of measles antibodies for at least two weeks.

If a patient is at risk of future measles exposure and receives a dose of less than 245 mg/kg subcutaneously per week, increase the dose to 245 mg/kg.

2.2 Preparation

CUTAQUIG is a clear and colorless solution that may turn to slightly opalescent and pale yellow during storage. Do not use the solution if it appears cloudy or contains particulates.

- Use aseptic technique when preparing and administering CUTAQUIG.
- Prior to administration visually inspect each vial of CUTAQUIG for particulate matter, whenever the solution and container permit.
- Do not mix CUTAQUIG with other products.
- Do not dilute CUTAQUIG.
- Do not shake the solution.
- Check the labeling for the expiration date and do not use beyond this date.
- CUTAQUIG vials are for single use only. Discard any unused product after each infusion in accordance with local requirements.
- Do not freeze. Do not use frozen product.
- Record lot number in patient's medical history.

2.3 Administration

CUTAQUIG is for subcutaneous use only.

Administer CUTAQUIG by a healthcare provider, or by caregiver or self-administered by the patient after appropriate training.

Infuse CUTAQUIG in the following areas: abdomen, thigh, upper arm, and/or upper leg/hip area.

CUTAQUIG is intended for subcutaneous use using an infusion pump and compatible syringe(s).

Infuse CUTAQUIG simultaneously into up to 6 different sites. Infusion sites should be at least 2 inches (5 cm) apart. Rotate infusion sites between subsequent administrations.

Volume:

For patients not already on IGSC therapy, do not exceed the maximum initial volume per injection site of 25 mL/site for adults (≥17 years), 15 mL/site (for ages 7-17 years) and 10 mL/site (for ages 2-6 years) for the first 2 infusions.

For subsequent infusions, if tolerated, you may gradually increase infusion volume/site by approximately 10 mL per infusion site in adults (≥17 years) and by approximately 5-10 mL per infusion site in children (2-17 years) every 2-4 weeks.

The maximum infusion volume is 40 mL/site (for adults ≥17 years), 29 mL/site (for ages 7-17 years) and 15.5 mL/site (for ages 2-6 years).

Rate:

For patients not already on IGSC therapy, do not exceed the maximum recommended flow rates per hour per infusion site of 20 mL per hour per site for adults (≥17 years) and 15 mL per hour per site for children (2-17 years) for the first 2 infusions.

For subsequent infusions, if tolerated, you may gradually increase the infusion rate by approximately 10 mL/hr/site in adults (≥17 years) and by approximately 5-10 mL/hr/site in children (2-17 years) every 2-4 weeks.

The maximum infusion rate is 52 mL per hour per site for adults (≥17 years) and 25 mL per hour per site for children (2-17 years).

Table 2 Infusion Volumes and Infusion Rates (Infusion rate and infusion volume per site may gradually be increased every 2-4 weeks as tolerated by patients)

Infusion parameters*		Infusion Number	
		First 2 infusions	Subsequent infusions
Volume (mL/site)	Adults ≥17 years	≤ 25	gradually increase by approximately 10 mL/site every 2-4 weeks up to a maximum of 40
	7-17 years	≤ 15	gradually increase by approximately 5-10 mL/site every 2-4 weeks up to a maximum of 29
	2-6 years	≤ 10	gradually increase by approximately 5-10 mL/site every 2-4 weeks up to a maximum of 15.5
Rate (mL/hr/site)	Adults ≥17 years	≤ 20	gradually increase by approximately 10 mL/hr/site every 2-4 weeks up to a maximum of 52
	2-17 years	≤ 15	gradually increase by approximately 10 mL/hr/site every 2-4 weeks up to a maximum of 25

* As tolerated

Administration/Handling instructions

CUTAQUIG is for subcutaneous use only.

Follow the administration guidance below and use aseptic technique when administering CUTAQUIG.



Figure 1

1. Getting ready for infusion

- Choose and prepare a clean work area.
- Gather your infusion equipment:
 - Infusion pump and compatible syringe(s).
 - Needle or needleless transfer device (for drawing up product from the vial).
 - Infusion set (varies according to manufacturer's instructions).
 - Infusion tubing and Y-connector (if required).
 - Ancillary supplies: disinfectant wipes, gauze or transparent dressing, tape and sharps container.
 - Patient's treatment diary/log book.
- Wash your hands thoroughly and let them dry (*Figure 1*).



Figure 2

2. Checking and opening the vials

- If taken out of the refrigerator, allow products to reach room temperature ($\leq 25^{\circ}\text{C}$ / 77°F).
- Inspect each vial carefully:
 - Check that labelled dose is correct and based on prescription.
 - Check that the expiration date has not been passed.
 - Check the appearance of the solution: do not use the solution if it is cloudy or contains particles.
 - Make sure the protective cap is not broken or missing.
- Remove the protective cap.
- Disinfect the rubber stopper by using a sterile wipe and allow it to dry (*Figure 2*).




Figure 3

3. Preparing and filling the syringe

- Open sterile syringe and needle or needleless transfer device.
 - If a needleless transfer device is used, follow the instructions of the device manufacturer.
 - If the transfer is done using needle and syringe follow the instructions below:
 - Attach the needle to the syringe with a screw action.
 - Draw back on the plunger to fill the syringe with air which should be roughly equal to the amount of solution needed from the vial.
 - Insert the needle into the center of the vial stopper and turn the vial upside down. To avoid foaming, ensure that the tip of the needle is not in the solution, and then inject the air.
 - Next, move the needle so that the tip is in the solution, and then slowly draw up the desired volume of CUTAQUIG (*Figure 3*).
 - Withdraw the needle from the vial.
- This procedure might be repeated if you need to use more than one vial.
- When finished, remove the needle and dispose of it into the sharps bin and proceed to the next step.




Figure 4

4. Preparing the infusion pump and tubing

- Follow the manufacturer's instructions for preparing the infusion pump.
- Prime the administration tubing according to manufacturer's instruction to eliminate all remaining air (*Figure 4*).
- Stop priming before fluid reaches the needle tip.




Figure 5

5. Preparing the infusion site(s) and inserting the infusion needle(s)

- CUTAQUIG can be infused in the following areas: abdomen, thigh, upper arm, and/or upper leg/hip area (*Figure 5*).
- The number and location of injection sites depends on the volume of the total dose. The infusion sites should be at least 2 inches apart. A maximum of 6 infusion sites can be used simultaneously.
- Rotate sites between infusions.
- Avoid inserting the needle into scars, tattoos or injured/inflamed skin areas and inspect skin for signs of infection.
- Clean your skin at your selected infusion site(s) with an antiseptic skin wipe starting at the center and working outward in a circular motion, and allow each site to dry before proceeding.
- Pinch the skin between your thumb and forefinger around the injection site (*Figure 6*) and insert the needle into the subcutaneous tissue (*Figure 7*). The angle of the needle will depend on the type of infusion set being used.



Figure 6




Figure 7




Figure 8

- Secure the needle in place by applying sterile gauze and tape or transparent dressing (*Figure 8*).

6. Checking the infusion

- Check needle placement by pulling back on the syringe plunger. There should not be any blood return in the tubing.
- If blood return is seen, remove needle and restart from step 5 with new tubing at a different location.

7. Starting the infusion

- Start the infusion. Follow infusion pump manufacturer's instructions.

8. Recording the infusion

- On each vial of CUTAQUIG, you will find a peel-off portion of the label with the batch number details. Stick this label in your patient's treatment diary or infusion log book. Record details of the dose, date, time, infusion site location and any infections, side effects or other comments.

9. After infusion is complete

- Gently remove the needle(s) and immediately place into the sharps bin.
- If necessary, press a small piece of gauze on the needle site and apply a dressing.
- Discard all used disposable supplies as well as any unused product and the empty vial(s) as recommended by your healthcare provider and according to local requirements.
- Clean and store the infusion pump according to the manufacturer's instructions.

For self-administration, provide the patient and caregiver with instructions and appropriate training for infusion at home or other appropriate setting. Verify that the patient or caregiver is capable of self-administration or administration, respectively, using appropriate technique.

3 DOSAGE FORMS AND STRENGTHS

CUTAQUIG is a solution containing 16.5% IgG (165 mg/mL).

4 CONTRAINDICATIONS

CUTAQUIG is contraindicated:

- In patients who have had an anaphylactic or severe systemic reaction to the subcutaneous administration of human immune globulin or to any of the components of CUTAQUIG such as Polysorbate 80.
- In IgA-deficient patients with antibodies against IgA and a history of hypersensitivity to human globulin treatment.

5 WARNINGS AND PRECAUTIONS

5.1 Hypersensitivity

Severe hypersensitivity reactions may occur with CUTAQUIG, even in patients who tolerated previous treatment with human immune globulin. If a hypersensitivity reaction occurs, discontinue the CUTAQUIG infusion immediately and initiate appropriate treatment. Have epinephrine immediately available to treat any severe acute hypersensitivity reaction. IgA-deficient patients with known anti-IgA antibodies have a higher risk of developing potentially severe hypersensitivity and/or anaphylactic reactions (including anaphylaxis and shock) with administration of CUTAQUIG. CUTAQUIG contains less than or equal to 0.6 mg of IgA /mL.

5.2 Thrombosis

Thrombosis may occur following treatment with immune globulin products, including CUTAQUIG. Risk factors may include: advanced age, prolonged immobilization, hypercoagulable conditions, history of venous or arterial thrombosis, use of estrogens, indwelling central vascular catheters, hyperviscosity, and cardiovascular risk factors. Thrombosis may occur in absence of known risk factors.

Consider baseline assessment of blood viscosity in patients at risk for hyperviscosity, such as those with cryoglobulins, fasting chylomicronemia/markedly high triglycerides, or monoclonal gammopathies. For patients at risk of thrombosis, administer CUTAQUIG at the minimum dose and infusion rate practicable. Ensure adequate hydration in patients before administration. Monitor for signs and symptoms of thrombosis and assess blood viscosity in patients at risk for hyperviscosity. [see [WARNING](#), [Patient Counseling Information](#) (17)]

5.3 Falsely Elevated Glucose Readings

Blood Glucose Testing: Some types of blood glucose testing systems (for example, those based on the glucose dehydrogenase pyrroloquinolinequinone (GDH-PQQ) or glucose-dye-oxidoreductase methods) falsely interpret the maltose contained in CUTAQUIG as glucose. This may potentially result in falsely elevated glucose readings and, consequently, in the inappropriate administration of insulin, resulting in life-threatening hypoglycemia. Also, cases of true hypoglycemia may go untreated if the hypoglycemic state is masked by falsely elevated glucose readings. Accordingly, when administering CUTAQUIG, measure blood glucose with a glucose-specific method. The product information of the blood glucose testing system, including that of the test strips, should be carefully reviewed to determine if the system is appropriate for use with maltose-containing parenteral products. If any uncertainty exists, contact the manufacturer of the testing system to determine if the system is appropriate for use with maltose-containing parenteral products.

5.4 Aseptic Meningitis Syndrome (AMS)

Aseptic Meningitis Syndrome can occur with CUTAQUIG. AMS has been reported after the use of human immune globulin administered intravenously and subcutaneously. It usually begins a few hours to 2 days following treatment and occurs more frequently in females than in males.

AMS is characterized by the following signs and symptoms: severe headache, nuchal rigidity, drowsiness, fever, photophobia, painful eye movements, nausea and vomiting. Cerebrospinal fluid (CSF) studies frequently show pleocytosis up to several thousand cells per mm³, predominantly from the granulocytic series, and elevated protein levels up to several hundred milligram/dL, but negative culture results. To rule out other causes of meningitis, conduct a thorough neurological examination on patients showing such symptoms and signs, including CSF studies. Discontinuation of immunoglobulin treatment has resulted in remission of AMS within several days without sequelae.

5.5 Renal Dysfunction/Failure

Acute renal dysfunction/failure, acute tubular necrosis, proximal tubular nephropathy, osmotic nephrosis and death may occur with use of human immune globulin, especially those containing sucrose. CUTAQUIG does not contain sucrose. Ensure that patients are not volume-depleted before administration of CUTAQUIG.

In patients at risk of developing renal dysfunction because of any degree of preexisting renal insufficiency or predisposition to acute renal failure (such as diabetes mellitus, age greater than 65 years, volume depletion, sepsis, paraproteinemia, or patients receiving known nephrotoxic drugs) monitor renal function and consider lower, more frequent dosing [see [Dosage and Administration \(2\)](#), [Patient Counseling Information \(17\)](#)].

Periodic monitoring of renal function and urine output is particularly important in patients judged to have a potential increased risk of developing acute renal failure. Assess renal function, including measurements of blood urea nitrogen (BUN) and serum creatinine, before the initial infusion of CUTAQUIG and again at appropriate intervals thereafter. If renal function deteriorates, consider discontinuing CUTAQUIG.

5.6 Hemolysis

IgG products, including CUTAQUIG, can contain blood group antibodies that may act as hemolysins and induce *in vivo* coating of red blood cells (RBCs) with immunoglobulin, causing a positive direct antiglobulin (Coombs') test result. Delayed hemolytic anemia can develop subsequent to immune globulin therapy due to enhanced RBC sequestration, and acute hemolysis, consistent with intravascular hemolysis, has been reported.

Monitor CUTAQUIG recipients for clinical signs and symptoms of hemolysis, particularly patients with risk factors (such as non-O blood group, administration of high IgG doses ($\geq 2\text{g/kg BW}$)). Underlying inflammatory state in a patient may increase the risk of hemolysis but its role is uncertain. Consider appropriate confirmatory laboratory testing if signs and symptoms of hemolysis are present after CUTAQUIG infusion.

5.7 Transfusion-related Acute Lung Injury (TRALI)

Non-cardiogenic pulmonary edema may occur in patients administered human immune globulin products. TRALI is characterized by severe respiratory distress, pulmonary edema, hypoxemia, normal left ventricular function, and fever. Typically, it occurs within 1 to 6 hours following transfusion.

Monitor patients for pulmonary adverse reactions. If TRALI is suspected, perform appropriate tests for the presence of anti-neutrophil antibodies in both the product and patient's serum. Patients with TRALI may be managed using oxygen therapy with adequate ventilatory support.

5.8 Transmittable Infectious Agents

Because CUTAQUIG is made from human plasma, it may carry a risk of transmitting infectious agents, e.g., viruses, the variant Creutzfeldt-Jakob disease (vCJD) agent and, theoretically, the Creutzfeldt-Jakob disease (CJD) agent. This also applies to unknown or emerging viruses and other pathogens. No cases of transmission of viral diseases or CJD have been associated with the use of CUTAQUIG. All infections suspected by a physician to have possibly been transmitted by CUTAQUIG should be reported to Pfizer Inc. at 1-800-438-1985.

5.9 Interference with Laboratory Tests

After infusion of CUTAQUIG, the transitory rise of the various passively transferred antibodies in the patient's blood may yield false positive serological test results, with the potential for misleading interpretation. Passive transmission of antibodies to erythrocyte antigens (e.g., A, B, and D) may cause a positive direct or indirect antiglobulin (Coombs') test.

6 ADVERSE REACTIONS

The most common adverse reactions (AR) observed in $\geq 5\%$ of study subjects were local infusion site adverse reactions (such as redness, swelling, and itching), headache, fever, dermatitis, asthma, diarrhea, and cough.

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

Clinical safety data are based on one pivotal open-label, single arm, prospective, multicenter major effectiveness study of CUTAQUIG in subjects with primary humoral immunodeficiency (PI), previously treated with Immune Globulin Intravenous (Human) (IGIV) for at least 6 months and its extension study. The pivotal study was conducted in Europe and North

America and was followed by an extension safety study (prospective, open-label, single-arm, multicenter phase 3). 21 patients were initially treated in the pivotal study and 6 patients from Canada were newly treated. Total number of patients in both studies combined was 81.

In the pivotal study, the 75 subjects in the Safety Analysis Set received 4462 infusions, with a mean of 59 infusions administered per subject. The average dose of CUTAQUIG used per subject was 0.187 g/kg in adult subjects, 0.150 g/kg in young children, 0.164 g/kg in older children and 0.170 g/kg in adolescents.

In the extension study, the 27 subjects in the Safety Analysis Set received 2777 infusions. The average dose of CUTAQUIG used per subject was 0.166 g/kg in adult subjects, 0.127 g/kg in young children, 0.210 g/kg in older children and 0.160 g/kg in adolescents.

Excluding infections, a total of 74 out of 81 subjects (91%) experienced 1527 adverse reactions (ARs, defined as adverse events occurring during or within 72 hours of infusion or any adverse events otherwise causally related) in both studies. Of the 1527 ARs 1088 were infusion site reactions. The proportion of infusions with adverse reactions was 0.21. Sixty-one subjects (75%) had at least one systemic AR.

Local reactions were the most common ARs and were experienced by 59 subjects (73%). The rate of infusion site reaction per infusion was 0.15. Most local ARs were either mild (transient AR causing discomfort but not interfering with routine activities [87.5%]) or moderate (AR sufficiently discomforting to interfere with routine activities [10.3%]) in intensity.

[Table 3](#) and [Table 4](#) summarize adverse reactions occurring in $\geq 5\%$ of adult subjects and pediatric subjects in both the pivotal and the extension study.

Table 3 Adverse reactions* in $\geq 5\%$ of adult subjects and rate per infusion in adult subjects (pivotal and extension study)

ARs	Number (%) of subjects (N=43)	Number (rate**) of ARs (N=3956 infusions)
Local reaction	31 (72.1)	648 (16.4)
Systemic ARs		
Headache	8 (18.6)	10 (0.002)
Dermatitis	5 (11.6)	6 (0.001)
Fever	5 (11.6)	5 (0.001)
Diarrhoea	5 (11.6)	7 (0.002)
Muscle spasms	4 (9.3)	5 (0.001)
Back pain	4 (9.3)	4 (0.001)
Arthralgia	3 (7.0)	3 (0.001)

* Excluding infections.

** Rate = total number of adverse reactions divided by total number of infusions.

Table 4 Adverse reactions* in $\geq 5\%$ pediatric subjects and rate per infusion in pediatric subjects (pivotal and extension study)

ARs	Number (%) of subjects (N=38)	Number (rate**) of ARs (N=3283 infusions)
Local reaction	28 (73.7)	442 (13.5)
Systemic ARs		
Asthma	4 (10.5)	7 (0.002)
Cough	4 (10.5)	5 (0.002)
Vomiting	4 (10.5)	5 (0.002)
Nasal congestion	3 (7.9)	4 (0.001)
Fever	3 (7.9)	4 (0.001)
Headache	3 (7.9)	3 (0.001)
Alanine aminotransferase increased	3 (7.9)	3 (0.001)
Leukopenia	3 (7.9)	3 (0.001)

ARs	Number (%) of subjects (N=38)	Number (rate**) of ARs (N=3283 infusions)
Neutropenia	3 (7.9)	3 (0.001)
Dermatitis	2 (5.3)	3 (0.001)
Oropharyngeal pain	2 (5.3)	3 (0.001)
Urticaria	2 (5.3)	2 (0.001)
Aspartate aminotransferase increased	2 (5.3)	2 (0.001)
Abdominal pain	2 (5.3)	2 (0.001)
Ear pain	2 (5.3)	2 (0.001)

* Excluding infections.

** Rate = total number of adverse reactions divided by total number of infusions.

The most frequent infusion site reactions were redness, swelling, and itching. During the pivotal study period the incidence of local reactions decreased over time from 37% of infusions triggering an infusion site reaction during the first 4 infusions to around 16% of infusions with an infusion site reaction during the last 4 infusions.

6.2 Postmarketing Experience

The following adverse reactions have been identified during post-approval use of immune globulin products administered subcutaneously. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

[Table 5](#) summarizes adverse reactions that have been reported during post-marketing use of immune globulin products:

Table 5 Adverse reactions reported during post-marketing use of immune globulin products

MedDRA System Organ Class (SOC)	Adverse reactions
Blood and lymphatic system disorders	Pancytopenia, leukopenia, hemolytic anemia
Immune system disorders	Anaphylactic reaction, hypersensitivity reaction, allergic reaction, angioneurotic edema, face edema
Psychiatric disorders	Agitation
Nervous system disorders	Loss of consciousness, cerebrovascular accident, aseptic meningitis, seizures, migraine, tremor, paresthesia, dizziness
Cardiac disorders	Cardiac arrest, tachycardia, palpitations
Vascular disorders	Thromboembolism, thrombosis, circulatory collapse, hypertension, hematoma
Respiratory, thoracic and mediastinal disorders	Transfusion-related acute lung injury, acute respiratory distress syndrome, respiratory failure, pulmonary embolism, apnea, cyanosis, hypoxia, pulmonary edema, bronchospasm, dyspnea, cough, wheezing
Gastrointestinal disorders	Hepatic function abnormal, nausea, vomiting, abdominal distension, abdominal pain upper
Skin and subcutaneous tissue disorders	Stevens-Johnson syndrome, epidermolysis, erythema multiforme, eczema, urticaria, rash (erythematous), alopecia, skin discoloration, skin mass, skin reaction, skin/infusion site ulceration, skin/infusion site necrosis
Musculoskeletal and connective tissue disorders	Back pain, arthralgia, pain in extremity, myalgia
Renal and urinary disorders	(Acute) renal failure
General disorders and administration site condition	Chills, chest pain, chest discomfort, hot flush, flushing, hyperhidrosis, fatigue, influenza-like illness, malaise
Investigations	Hepatic enzyme increased, coombs test positive, free hemoglobin present, hemoglobin increased, haptoglobin decreased

The following adverse reactions have been identified during post-approval use of CUTAQUIG: dizziness, nausea, pruritus and fatigue. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

To report SUSPECTED ADVERSE REACTIONS, contact Octapharma USA Inc. at 1-866-766-4860 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

7 DRUG INTERACTIONS

7.1 Serological Testing

Various passively transferred antibodies in immunoglobulin preparations may lead to misinterpretation of results of serological testing.

7.2 Live Virus Vaccines

The passive transfer of antibodies with immunoglobulin administration may interfere with the response to live virus vaccines such as measles, mumps, rubella, and varicella. Inform the immunizing medical practitioner of recent therapy with CUTAQUIG so that appropriate precautions can be taken. [See [Patient Counselling Information \(17\)](#)]

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

No human data are available to indicate the presence or absence of drug-associated risk. Animal reproduction studies have not been conducted with CUTAQUIG. It is not known whether CUTAQUIG can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Immune globulins cross the placenta from maternal circulation increasingly after 30 weeks of gestation. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2-4% and 15-20%, respectively.

8.2 Lactation

Risk summary

No human data are available to indicate the presence or absence of drug associated risk. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for CUTAQUIG and any potential adverse effects on the breastfed infant from CUTAQUIG or from the underlying maternal condition.

8.4 Pediatric Use

The safety and efficacy of CUTAQUIG have been established for primary humoral immunodeficiency in the pediatric age group 2 to 17 years. CUTAQUIG was evaluated in 38 pediatric subjects with primary humoral immunodeficiency (26 children and 12 adolescents) in the pivotal and extension studies. [see [Clinical Studies \(14\)](#)]

Overall safety and efficacy findings in pediatric age group were comparable to those seen in adults. However, infusion site reactions were more commonly observed in adults compared to the pediatric population.

PK parameters are comparable between pediatric and adult patients. No pediatric-specific dose requirements were necessary to achieve the desired serum IgG levels.

Safety and effectiveness of CUTAQUIG in pediatric patients below the age of 2 have not been established.

8.5 Geriatric Use

Clinical studies of CUTAQUIG did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Three study subjects enrolled in the clinical trial were 65 years and over. In general, dose selection for an elderly patient should be cautious. Medical practitioners should start at the low end of the dosing range to reduce the risk of hyperviscosity and precipitation of cardiac, renal, or hepatic adverse reaction and of concomitant disease or other drug therapy.

11 DESCRIPTION

CUTAQUIG (Immune Globulin Subcutaneous (Human) - hipp), is a solvent/detergent (S/D)-treated, sterile preparation of highly purified immunoglobulin G (IgG) derived from large pools of human plasma. CUTAQUIG is a solution for injection to be administered subcutaneously.

This preparation contains approximately 165 mg of protein per mL (16.5%), of which not less than 96% is normal human immunoglobulin G. CUTAQUIG contains not more than 3% aggregates, not less than 94% monomers and dimers, and not more than 3% fragments. The product contains equal to or less than 0.6 mg of IgA /mL.

The sodium content of the final solution is not more than 30 mmol/L and the pH is between 5.0 and 5.5. The osmolality is 310 - 380 mOsmol/kg.

The manufacturing process for CUTAQUIG isolates IgG without additional chemical or enzymatic modification, and the Fc portion is maintained intact. CUTAQUIG contains the IgG antibody activities present in the donor population. IgG subclasses are fully represented with the following approximate percent of total IgG: IgG1 is 70%, IgG2 is 25%, IgG3 is 3% and IgG4 is 2%.

CUTAQUIG contains a broad spectrum of IgG antibodies against bacterial and viral agents that are capable of opsonization and neutralization of microbes and toxins. It contains maltose (79 mg/mL), but no preservatives or sucrose.

All units of human plasma used in the manufacture of CUTAQUIG are provided by FDA-approved blood and plasma establishments, and are tested by FDA-licensed serological tests for HBsAg, antibodies to HCV and HIV and Nucleic Acid Test (NAT) for HCV and HIV-1 and found to be non-reactive (negative).

The product is manufactured by the cold ethanol fractionation process followed by ultrafiltration and chromatography. The manufacturing process includes treatment with an organic S/D mixture composed of tri-n-butyl phosphate (TNBP) and Octoxynol. The CUTAQUIG manufacturing process shows significant viral reduction in in vitro studies (Table 6). These reductions are achieved through a combination of process steps including cold ethanol fractionation, S/D treatment and pH 4 treatment.

Table 6 Pathogen Reduction During CUTAQUIG Manufacturing

Production Step	In Vitro Reduction Factor [\log_{10}]				
	Enveloped Viruses			Non-Enveloped Viruses	
	HIV-1	PRV	SBV	MEV	PPV
Ethanol fractionation	≥ 4.81	≥ 6.28	≥ 7.13	≥ 7.13	≥ 6.53
S/D treatment	≥ 4.93	5.23	≥ 6.77	n.a.	
pH4 treatment	≥ 4.33	≥ 6.71	6.71	5.07	$< 1^*$
Global reduction factor	≥ 14.07	≥ 18.22	≥ 20.61	≥ 12.20	≥ 6.53

* Not calculated for global \log_{10} Reduction Factor

HIV-1: Human Immunodeficiency Virus - 1

PRV: Pseudorabies Virus, model virus for e.g. Hepatitis B Virus (HBV)

SBV: Sindbis Virus, model virus for Hepatitis C Virus (HCV)

MEV: Mouse Encephalomyelitis Virus, model virus for Human Parvovirus B19

PPV: Porcine Parvovirus, model virus for Hepatitis A Virus (HAV)

n.a.: not applicable

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

CUTAQUIG supplies a broad spectrum of opsonizing and neutralizing Immunoglobulin G (IgG) antibodies against a wide variety of bacterial and viral agents. It has a distribution of immune globulin subclasses closely proportional to that in native human plasma. The mechanism of action in primary humoral immunodeficiency (PI) has not been fully elucidated; however adequate doses may restore abnormally low immune globulin G levels to the normal range and thus help in preventing infections.

12.2 Pharmacodynamics

CUTAQUIG contains mainly IgG with a broad spectrum of antibodies against various infectious agents reflecting the IgG activity found in the donor population. CUTAQUIG which is prepared from pooled material from not less than 1000 donors, has an IgG subclass distribution similar to that of native human plasma. Adequate doses of Immune Globulin Subcutaneous (Human) (IGSC) can restore an abnormally low IgG level to the normal range. Standard pharmacodynamic studies were not performed.

12.3 Pharmacokinetics

A pharmacokinetic (PK) sub-study was conducted in 37 subjects (18 adults and 19 pediatric subjects) who were enrolled in the pivotal safety and efficacy study. Blood samples for PK study were collected prior to switching to CUTAQUIG (IGIV profile: PK_{IV}), after the 11th infusion of CUTAQUIG (first SC profile: PK_{SC1}) and after the 28th infusion of CUTAQUIG (second SC profile: PK_{SC2}). The objective of the PK sub-study was to compare the AUCs following the IV and SC administration. At steady state, the ratio of geometric least square means (SC2:IV) was 1.06, (90% CI: 1.03, 1.10), indicating comparable exposure between IGSC and IGIV treatment (standardized to a 7-day period). The actual dose converting factor was 1.41 (1.21, 1.89). Using a Population PK model calculation, the DCF was determined in a statistically more advanced manner at 1.33 for a median patient.

Serum IgG and IgG subclass trough levels were nearly constant during the IGSC phase of the study, with higher mean levels after SC treatment compared with those following IGIV. At the end of the IGIV period, trough levels ranged from 5.0 g/L to 15.1 g/L. Over the entire IGSC treatment period individual trough levels of total IgG ranged between 4.4 g/L to 24.0 g/L.

[Table 7](#) summarizes the Key PK parameters for CUTAQUIG. PK parameters are comparable between pediatric and adult patients.

Table 7 Key Pharmacokinetic Parameters for CUTAQUIG and IGIV

Parameter [Arithmetic Mean (SD)]	IGIV (N=37)	CUTAQUIG (N=37)
C _{max} [g/L]	18.01 (4.5)	13.47 (3.7)
C _{min} [g/L]	10.09 (2.5)	11.66 (2.9)
T _{max} [h] #	3.38 (1.6 – 69.5)	49.62 (0.8 – 98.3)
AUC _{tau} [g*hr/L]	2013 (570)*	2233 (586)
AUC _{tau} [mg*day/dL]	8389 (2376)*	9302 (2441)
CL ⁺ (mL/day/kg)	1.5 (0.5)	1.9 (0.6)
Actual IgG Dose per kg Body Weight and Week (g/kg/week)	0.121 (0.049)	0.170 (0.066)

T_{max} is presented as Median (range)

* standardized to a 7-day period

+ apparent clearance (CL/F) for CUTAQUIG (F = bioavailability)

Compared with weekly administration, administration of CUTAQUIG on a every other week basis at double the weekly dose is expected to result in comparable IgG exposure [equivalent AUCs, with a slightly higher IgG peak (C_{max}) and slightly lower trough (C_{min})] in both adult and pediatric subjects. In addition, for the same total weekly dose, CUTAQUIG infusions given 2, 3, 5, or 7 times per week (frequent dosing) are expected to produce IgG exposures comparable to weekly dosing in both adult and pediatric subjects.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

No animal studies were conducted with CUTAQUIG to assess carcinogenesis, mutagenesis or impairment of fertility.

13.2 Animal Toxicology and/or Pharmacology

Safety of CUTAQUIG has been demonstrated in several standard nonclinical toxicology studies (local tolerance in rabbits, cardiovascular and respiratory effects in dogs, thrombogenic potential in rabbits).

The product given subcutaneously to CD1 mice inoculated with *Streptococcus pneumoniae* showed a dose related improvement in survival.

TNBP and Octoxynol may be found in CUTAQUIG in trace amounts. In animals and in vitro studies, these compounds were not genotoxic and did not have mutagenic properties. They did not show teratogenic effects when administered to pregnant rabbits and rats during organogenesis.

14 CLINICAL STUDIES

There were 2 clinical studies, one was a pivotal, prospective, open-label, single-arm, multicenter study to evaluate the pharmacokinetics (PK), efficacy, tolerability and safety of subcutaneous human immunoglobulin (CUTAQUIG) in subjects with primary humoral immunodeficiency (PI). The other was an extension study.

The pivotal study was conducted in 75 subjects (37 adult and 38 pediatric subjects < 17 years of age) who received weekly SC infusions with CUTAQUIG during a 12-week wash-in/wash-out period followed by a 12-month efficacy period during which efficacy, pharmacokinetics, safety, tolerability, and quality of life (QoL) parameters of CUTAQUIG were evaluated. All patients received study treatment and 68 patients completed the study. Seven patients (4 adolescents and 3 adults) were withdrawn prematurely. Reasons for withdrawal from the study were patient's decision in 6 cases and patient's noncompliance in one adolescent patient.

During the efficacy period the mean weekly dose was 174 mg/kg BW, with individual doses ranging from 60 to 390 mg/kg BW. The median duration of infusion per week was 1.5 hours.

All enrolled subjects (n=75) were included in the Safety Analysis Set and the Full Analysis Set (FAS). Four subjects were excluded from the Per-Protocol (PP) Set because they terminated early before the start of the primary treatment period.

Overall, 36 female subjects and 39 male subjects participated in this study. The youngest subject enrolled in the study was 2 years old and the oldest was 73 years old. The mean age in the adult group (17–75 yrs) was 47.5 years. The mean age at time of enrollment in the pediatric groups was 4.2 years, 7.9 years and 14.1 years in the 3 pediatric age groups [young children (ages 2-5 years), older children (ages 6-11 years) and adolescents (ages 12-16 years)] respectively. Reported race was white for all but one subject, and all subjects were of not Hispanic/Latino ethnicity.

The main objective of the pivotal study was to assess the efficacy of CUTAQUIG in preventing serious bacterial infections (SBI defined as bacteremia/sepsis, bacterial meningitis, osteomyelitis/septic arthritis, bacterial pneumonia and visceral abscess). This endpoint was considered successful if the upper bound of the one-sided 99% confidence interval for the rate of SBIs was < 1.0 per subject-year of follow up. This criterion was met, as no SBIs were reported at any time during the study.

Other endpoints of the pivotal study included, but were not limited to: the number of episodes of any other infections, along with type and severity of infection and time to resolution; number of days of use and annual rate of antibiotics; absence and number of days of absence from work/school/ kindergarten/day care; and hospitalizations due to infections and number of days and annual rate of hospitalisation.

Efficacy results of the pivotal study are summarized in [Table 8](#).

Table 8 Summary of efficacy results of the pivotal study (FAS).

	Adults	Pediatrics	Total
Number of subjects (efficacy period)	37	38	75
Total number of subject years	35.5	35.0	70.5
Infections			
Annual rate [number of SBIs* per subject-year (upper one-sided 99% CI)]	0 (0.13) [#]	0 (0.13) [#]	0 (0.06) [#]
Annual rate [number of other infections per subject-year (two-sided 95% CI)]	3.4 (2.2, 5.4)	3.1 (2.0, 4.8)	3.3 (2.4, 4.5)
Systemic antibiotic use			
Number of subjects (%)	25 (67.6%)	24 (63.2%)	49 (65.3%)
Annual rate [days on treatment per subject-year (two-sided 95%CI)]	32.1 (17.2, 60.1)	62.6 (31.0, 126.4)	47.2 (28.4, 78.6)

	Adults	Pediatrics	Total
Number of subjects (efficacy period)	37	38	75
Days out of work/school/kindergarten/day care due to infections			
Number of days	72	180	252
Annual rate [days per subject-year (two-sided 95% CI)]	2.0 (0.7, 5.7)	5.2 (2.8, 9.4)	3.6 (2.1, 6.0)
Hospitalization due to infections	0	4	4
Number of days	0	29	29
Annual rate [days per subject-year (two-sided 95% CI)]	0 (0.1) [#]	0.8 (0.3, 2.6)	0.4 (0.1, 1.3)

* Defined as bacterial pneumonia, bacteremia/septicemia, osteomyelitis/septic arthritis, bacterial meningitis, and visceral abscess.

CI = confidence interval

[#] calculation for the upper limit of two-sided 95% or one-sided 99% CI was based on standard Poisson model for zero counts

The extension study was a prospective, open-label, single-arm, multicenter phase 3 safety study that enrolled 27 patients (17 adults, 10 patients aged <17 years) with PI. Twenty-one patients were initially treated in the pivotal study and 6 patients were newly enrolled. Mean age was 39 years (range 6 to 73 years). Ten patients (37%) were male, all but 2 were White, none was Hispanic or Latino.

Patients received CUTAQUIG on a weekly (25 patients) or an “every other week” schedule (2 patients). The primary objective of this study was to assess the medium-to-long term safety and tolerability of CUTAQUIG. Secondary efficacy assessments included occurrence of SBIs, the annual rate of all infections of any kind or seriousness, hospitalizations due to infections, and antibiotic use. One adult had bacteremia/sepsis. The rate of SBI per person-year was 0.03 for adults, and 0.0 for all other age groups (overall rate of 0.018). Secondary efficacy results are summarized in [Table 9](#).

Table 9 Summary of efficacy results of the extension study (FAS)

	Adults	Pediatrics	Total
Number of subjects	17	10	27
Total number of subject years	33.4	20.7	54.1
Infections			
Annual rate [number of SBIs* per subject year (upper limit of one-sided 99% CI)]	0.03 (0.31)	0 (0.22) [#]	0.02 (0.19)
Annual rate [number of other infections per subject-year (two-sided 95% CI)]	2.3 (1.0, 4.9)	2.0 (0.9, 4.8)	2.2 (1.2, 3.9)
Systemic antibiotic use			
Number of subjects (%)	13 (76.5%)	6 (60.0%)	19 (70.4%)
Annual rate [treatment days per subject-year (two-sided 95% CI)]	30.9 (12.2, 78.4)	70.6 (22.7, 220.0)	46.0 (21.3, 99.4)
Days out of work/school/kindergarten/day care due to infections			
Number of days	55	75	130
Annual rate [days per subject-year (two-sided 95% CI)]	1.7 (0.4 to 6.5)	3.6 (1.3 to 10.3)	2.4 (1.0, 5.5)
Hospitalization due to infections	3	0	3
Number of days	10	0	10
Annual rate [days per subject-year (two-sided 95% CI)]	0.3 (0.07 to 1.3)	0 (0.2) [#]	0.2 (0.04, 0.8)

* Defined as bacterial pneumonia, bacteremia/septicemia, osteomyelitis/septic arthritis, bacterial meningitis, and visceral abscess.

CI = confidence interval

[#] calculation for the upper limit of two-sided 95% or one-sided 99% CI was based on standard Poisson model for zero counts

16 HOW SUPPLIED/STORAGE AND HANDLING

CUTAQUIG is supplied in 1 g, 1.65 g, 2 g, 3.3 g, 4 g or 8 g single-use vials.

Carton NDC Number	Container NDC Number	Size	Grams Protein
0069-1061-02	0069-1061-01	6 mL	1
0069-1802-02	0069-1802-01	10 mL	1.65
0069-1476-02	0069-1476-01	12 mL	2
0069-1960-02	0069-1960-01	20 mL	3.3
0069-1509-02	0069-1509-01	24 mL	4
0069-1965-02	0069-1965-01	48 mL	8

- Components used in the packaging of CUTAQUIG are not made with natural rubber latex.
- Store at +2°C to +8°C (36°F to 46°F) for up to 36 months from the date of manufacture. Within its shelf-life, the product may be stored at room temperature up to +25°C (77°F) for up to 6 months without being refrigerated again during this period, and must be discarded if not used after this.
- Keep the vial in the outer carton to protect it from light.
- Check the product expiration date on the vial label. Do not use beyond the expiration date.
- Do not freeze. Do not use frozen product.
- Do not mix with other products.
- Do not shake the vial.
- Use aseptic technique when preparing and administering CUTAQUIG. Prior to administration, visually inspect each vial for particulate matter, whenever the solution and container permit. Do not use if the solution is cloudy or contains particulates.
- The vial is for single-use only. Discard any unused product after each infusion, in accordance with local requirements.

17 PATIENT COUNSELING INFORMATION

Advise the patients to read the FDA-approved patient labeling (Patient Information and Instructions for Use).

Inform patients to immediately report the following signs and symptoms to their healthcare provider:

- Hypersensitivity reactions to CUTAQUIG (including hives, generalized hives, tightness of the chest, wheezing, low blood pressure, and anaphylaxis): Advise patients to have epinephrine immediately available to treat any acute severe hypersensitivity reaction and ensure patients and their caregivers are properly trained in its use. [see [Warnings and Precautions \(5.1\)](#)].
- Pain and/or swelling of an arm or leg with warmth over the affected area, discoloration of an arm or leg, unexplained shortness of breath, chest pain or discomfort that worsens on deep breathing, unexplained rapid pulse, or numbness or weakness on one side of the body [see [Warnings and Precautions \(5.2\)](#)].
- Severe headache, neck stiffness, drowsiness, fever, sensitivity to light, painful eye movements, nausea, and vomiting [see [Warnings and Precautions \(5.4\)](#)].
- Decreased urine output, sudden weight gain, fluid retention/edema, and/or shortness of breath [see [Warnings and Precautions \(5.5\)](#)].
- Fatigue, increased heart rate, yellowing of the skin or eyes, and dark-colored urine [see [Warnings and Precautions \(5.6\)](#)].
- Severe breathing problems, lightheadedness, drops in blood pressure, and fever [see [Warnings and Precautions \(5.7\)](#)].

Inform patients to be tested periodically to make sure they have the appropriate levels of IgG in their blood. These tests may result in adjustments to the CUTAQUIG dose.

Inform patients/caregivers that because CUTAQUIG is made from human blood, it may carry a risk of transmitting infectious agents, e.g., viruses, the variant Creutzfeldt-Jakob disease (vCJD) agent and, theoretically, the Creutzfeldt-Jakob disease (CJD) agent [see [Warnings and Precautions \(5.8\)](#), [Description \(11\)](#)].

Inform patients that CUTAQUIG may interfere with the response to live virus vaccines (e.g., measles, mumps, rubella, and varicella) and to notify their immunizing physician of recent therapy with CUTAQUIG [see [Drug Interactions \(7.2\)](#)].

Home treatment/Self administration

- If self-administration is deemed appropriate by the physician, it is highly important to clearly instruct the patient/caregiver and to provide a thorough training on subcutaneous administration of CUTAQUIG. The patient/caregiver needs to demonstrate the ability to independently administer subcutaneous infusions and document it.
- Ensure the patient/caregiver understands the importance of regular subcutaneous infusions to maintain appropriate steady state IgG levels.
- Instruct the patient/caregiver to keep a treatment diary/log where all information about each infusion should be clearly documented.
- Inform patients to interrupt or terminate the CUTAQUIG infusion if a hypersensitivity reaction occurs.
- Inform patients that they should be tested regularly to make sure they have the correct levels of CUTAQUIG (IgG) in their blood. These tests may result in adjustment to the CUTAQUIG dose.
- Instruct the patient/caregiver that local injection site reactions (such as swelling, redness, itching) are a common side effect of subcutaneous treatment, but also tell them to contact their healthcare provider if local reactions increase in severity or persist for more than a few days.
- Inform the patient that injection sites should be rotated between infusions and that CUTAQUIG is for subcutaneous infusion only.

Manufactured by:
Octapharma Pharmazeutika Produktionsges.m.b.H.
Oberlaaer Strasse 235
1100 Vienna, Austria
U.S. License No. 1646

Distributed by:
Pfizer Labs
Division of Pfizer Inc.
NY, NY 10017

PATIENT PACKAGE INSERT

Patient Information

CUTAQUIG

(kew' ta kwig)

Immune Globulin Subcutaneous (Human) - hipp, 16.5% solution

Information for Patients

The following summarizes important information about CUTAQUIG. Please read it carefully before using CUTAQUIG and each time you get a refill, as there may be new information. This Patient Information does not take the place of talking with your healthcare provider about your medical condition or your treatment. If you have any questions after reading this, ask your healthcare provider.

What is CUTAQUIG?

CUTAQUIG is a ready-to-use liquid solution of Immunoglobulin G (IgG), also called antibodies, which protect the body against infection. CUTAQUIG is used to treat patients with primary humoral immunodeficiency (PI).

There are many forms of PI. The most common types of PI result in an inability to make a very important type of protein called antibodies, which help the body fight off infections from bacteria or viruses. Regular administration of CUTAQUIG will help your body to fight bacteria and viruses that cause infections. CUTAQUIG is made from human plasma that is donated by healthy people. CUTAQUIG contains antibodies collected from these healthy people; these antibodies replace the missing antibodies in PI patients.

Who should NOT use CUTAQUIG?

Do not use CUTAQUIG if you have ever had a severe allergic reaction to immune globulin or other blood products.

Tell your healthcare provider if you:

- ever had any severe reaction to other immune globulin medicines.
- were told that you have a condition called IgA deficiency.
- have a history of heart or blood vessel disease.
- have had blood clots or “thick blood”.
- have been immobile for some time.

What should I tell my healthcare provider before using CUTAQUIG?

Talk to your healthcare provider about any medical conditions that you have or have had.

Tell your healthcare provider that you are taking CUTAQUIG before you get vaccination as vaccines may not work while you are taking CUTAQUIG.

Tell your healthcare provider about all of the prescription and non-prescription medicines you take, including over-the-counter medicines, dietary supplements, or herbal medicines.

Tell your healthcare provider if you are pregnant or plan to get pregnant, or if you are nursing because CUTAQUIG might not be right for you.

Tell your healthcare provider if you have diabetes. If you need to do glucose testing, your healthcare provider may tell you to use a different way to monitor your blood sugar levels on the day that you receive a CUTAQUIG infusion. Some types of blood glucose testing systems (so called glucometers) falsely interpret the maltose contained in CUTAQUIG as glucose. If you are uncertain ask your healthcare provider which glucose testing system you can use while using CUTAQUIG.

How should I use CUTAQUIG?

CUTAQUIG is given under the skin (subcutaneously). This type of infusion can be given at home by yourself or by your caretaker after appropriate training. Administration of CUTAQUIG should be done at regular intervals. Only use CUTAQUIG by yourself after you have been instructed by your healthcare provider.

What are the possible or reasonably likely side effects of CUTAQUIG?

The most common side effects of CUTAQUIG are:

- Infusion site reactions (including but not limited to redness, swelling, itching, fluid in tissue, pain, mass, bruising)
- Headache
- Elevated body temperature

Call your healthcare provider or emergency department immediately if you have any of the following symptoms: difficulty breathing, chest tightness, itching, swelling of the face, rash, hives or dizziness. These could be signs of a serious allergic reaction.

Tell your healthcare provider immediately if you have any of the following symptoms. They could be signs of a serious drug reaction:

- Decreased urination, sudden weight gain, or swelling in the legs. These could be signs of a kidney problem.
- Bad headache with nausea, vomiting, stiff neck, fever, and sensitivity to light. These could be signs of meningitis, which is irritation of the lining around your brain.
- Pain, swelling, warmth, discoloration of an arm or leg, unexplained shortness of breath, chest pain, discomfort that worsens on deep breathing, unexplained rapid pulse, a lump in your legs or arms, or numbness or weakness on one side of the body. These could be signs of a blood clot.
- Chest pain or trouble breathing, or blue lips or extremities. These could be signs of a serious heart or blood problem.
- Brown or red urine, fast heart rate, yellow skin or eyes. These could be signs of a liver or heart problem.
- Fever over 100°F (38°C). This could be sign of an infection.

These are not all of the possible side effects from CUTAQUIG. Ask your healthcare provider for more information. You are encouraged to report side effects to Pfizer Inc. at 1-800-438-1985 or FDA at 1-800-FDA-1088.

Talk to your healthcare provider about any side effect that bothers you or that does not go away. You can ask your healthcare provider for more information on possible side effects.

Whenever giving yourself treatments at home, make sure another responsible person is present to help treat side effects or help if a serious adverse reaction occurs. Ask your healthcare provider whether you should have rescue medications, such as antihistamines or epinephrine. If prescribed epinephrine for severe allergic reactions, make sure you receive adequate training from a healthcare provider for its proper use.

How should I store CUTAQUIG?

Keep CUTAQUIG in the outer carton to protect it from exposure to light.

Do not freeze CUTAQUIG.

Store CUTAQUIG at 36°F- 46°F (+2°C to +8°C) for up to 36 months from the date of manufacture. Within its shelf-life, the product can be stored at room temperature up to 77°F (up to +25°C) for up to 6 months without being refrigerated again during this period, and must be discarded if not used after this.

Do not use CUTAQUIG after the expiration date printed on the vial.

Dispose all materials, including any unused CUTAQUIG, in an appropriate container.

What else should I know about CUTAQUIG?

Do not use CUTAQUIG for a medical condition for which it was not prescribed. Do not share CUTAQUIG with other people, even if they have the same diagnosis and symptoms that you have.

Manufactured by

Octapharma Pharmazeutika Produktionsges.m.b.H.
Oberlaaer Strasse 235
1100 Vienna, Austria

U.S. License No. 1646

Distributed by

Pfizer Labs

Division of Pfizer Inc.

NY, NY 10017

CUTAQUIG® is a registered trademark of Octapharma.

Patient Instructions for Use
CUTAQUIG
Immune Globulin Subcutaneous (Human) - hipp, 16.5% solution

Detailed patient handling instructions for administration of CUTAQUIG

CUTAQUIG is for subcutaneous use only.

Use CUTAQUIG only after you have been properly instructed and trained by your healthcare provider.

Follow the administration guidance below step by step and use aseptic/sterile technique when administering CUTAQUIG. Use gloves if you have been told to do so when preparing the infusion.

1. Prepare the necessary number of CUTAQUIG vials

- If stored in the refrigerator, put the vials at room temperature at least 90 minutes prior to infusion.
- Do not heat the vials or put them into the microwave.
- Do not shake the vials to avoid foaming.



Figure 1

2. Getting ready for infusion

- Choose and prepare a clean work area using antiseptic wipes or disinfecting solution.
- Gather your infusion equipment:
 - Infusion pump and compatible syringe(s)
 - Needle or needleless transfer device (for drawing up product from the vial)
 - Infusion set (varies according to manufacturer's instructions)
 - Infusion tubing and Y-connector (if required)
 - Ancillary supplies: disinfectant wipes, gauze or transparent dressing, tape and sharps container
 - Treatment diary and pen

- Wash and clean your hands thoroughly and let them dry as has been shown to you during the training (*Figure 1*). You can wear gloves during the preparation of the infusion if you have been told so during the training.

- If necessary, program the pump according to the user manual and as you have been shown during the training by your healthcare provider.

3. Checking and opening the vials

- Allow products to reach room temperature (77°F / ≤ 25°C).
- Inspect each vial carefully:

Check that the labelled dose is correct and based on your prescription.

Check that the expiration date has not been passed.

Check the appearance of the solution (it should be clear and colorless to pale yellow or slightly opalescent).



Figure 2

Do not use the solution if it is cloudy or contains particles.

Make sure the protective cap is not broken or missing.

- Remove the protective cap.
- Disinfect the rubber stopper by using a sterile wipe and allow it to dry (*Figure 2*).




Figure 3

4. Preparing and filling the syringe

- Open sterile syringe and needle or needleless transfer device.
- If a needleless transfer device is used, follow the instructions of the device manufacturer.
- If the transfer is done using needle and syringe, follow the instructions below:

Attach the needle to the syringe with a screw action.

Draw back on the plunger to fill the syringe with air. The amount of air should be roughly equal to the amount of solution needed from the vial.

Insert the needle into the center of the vial stopper and slowly turn the vial upside down. To avoid foaming, ensure that the tip of the needle is not in the solution; then inject air by pushing the plunger of the syringe.

Next, move the needle so that the tip is in the solution; then slowly draw up the desired volume of CUTAQUIG solution, making sure that the needle tip is always in the solution (*Figure 3*).

Withdraw the needle from the vial.

This procedure might be repeated if you need multiple vials for the calculated dose.

When finished, remove the needle and dispose of it into the sharps bin.

Once finished, proceed to the next step.




Figure 4

5. Preparing the infusion pump and tubing

- Follow the manufacturer's instructions for preparing the infusion pump.
- To prime the administration tubing, attach the filled syringe to the infusion tubing and gently push the plunger to fill the tubing with CUTAQUIG, as has been shown to you during training (*Figure 4*).
- Stop priming before fluid reaches the needle tip.




Figure 5

6. Preparing infusion site(s) and inserting the infusion needle(s)

- CUTAQUIG can be infused into the following areas: abdomen, thigh, upper arm, and/or upper leg/hip area (*Figure 5*).
- The number and location of injection sites depends on the volume of the total dose.
- The infusion sites should be at least 2 inches apart. Do not use more than 6 infusion sites at the same time.
- Rotate sites between infusions.
- Avoid inserting the needle into scars, tattoos, stretch marks or any skin that has signs of infection (such as injured/inflamed/red skin area).
- Clean your skin at your selected infusion site(s) with an antiseptic skin wipe, starting at the center and working outward in a circular motion. Allow each site to dry before proceeding.
- Pinch the skin between your thumb and forefinger around the injection site (*Figure 6*). Carefully remove the needle cover and insert the needle into the skin (*Figure 7*). The angle of the needle will depend on the type of infusion set being used.
- Put transparent dressing or sterile tape and gauze over the administration site to keep the needle in place during the infusion.




Figure 6




Figure 7




Figure 8

7. Checking the infusion

- Check needle placement by pulling back on the syringe plunger. There should not be any blood return in the tubing (*Figure 8*).
- If blood return is seen, remove the needle and restart from step 6 with new tubing at a different location.

8. Starting the infusion

- Start the infusion. Follow infusion pump manufacturer's instructions.

9. Recording the infusion

- On each vial of CUTAQUIG, you will find a peel-off portion of the label with the batch number details. Stick this label in your patient's treatment diary or infusion log book. Record details of the dose, date, time, infusion site location and any infections, side effects or other comments in connection with this infusion.

10. After infusion is complete

- Gently remove the dressing and the needle(s) and immediately place into the sharps bin.
- Press a small piece of gauze on the needle site and apply a dressing.
- Discard all used disposable supplies as well as any unused product and the empty vial(s) as recommended by your healthcare provider and according to local requirements.
- Tidy up and securely store all the reusable equipment (e.g., pump) until the next infusion.

If you encounter any problems or experience side effects during or after the infusion, contact your healthcare provider. When doing so, keep your treatment diary or log book with you to be able to give all necessary information. You can also report side effects to the FDA at 1-800-FDA-1088 or online under www.fda.gov/medwatch.

Manufactured by
Octapharma Pharmazeutika Produktionsges.m.b.H.
Oberlaaer Strasse 235
1100 Vienna, Austria
U.S. License No. 1646

Distributed by
Pfizer Labs
Division of Pfizer Inc.
NY, NY 10017

CUTAQUIG® is a registered trademark of Octapharma.

Issued 10 2021.