

Best Pharmaceuticals for Children Act (BPCA) Written Requests for Oncology Products

Denise Casey, MD

March 15, 2019



Goals of a Pediatric Development Program

- Obtain evidence based data to support the dosing, safety, and efficacy of products intended for use in the pediatric population
- Communicate this information in the product labeling to allow judicious use of the drug in the indicated population

Pediatric product development should be integrated into the adult development program and not be an afterthought.

1902 Biologics Control Act 1906 Pure Food and Drug Act 1938 Food Drug and Cosmetic Act 1962 Kefauver-Harris Amendment

Early pediatric legislation reflected a response to products that caused harm.



Later pediatric legislation encourages pediatric investigations to inform product labeling.

1979 Pediatric Use Subsection under Precautions

1997 FDAMA/Pediatric exclusivity provision

2002 Best Pharmaceuticals for Children Act (BPCA)

2003 Pediatric Research Equity Act (PREA)

2007 Food & Drug Administration Amendments Act (FDAAA)

2012 FDA Safety and Innovation Act (FDASIA)

2017 FDA Reauthorization Act (FDARA)

 PREA amended to require pediatric assessments of drugs directed at molecular targets considered relevant in pediatric cancer regardless of the adult indication; removes orphan exemption

Best Pharmaceuticals for Children Act (BPCA)



- Provides a financial incentive to <u>VOLUNTARILY</u> conduct pediatric studies of a product under a Written Request (WR)
- A WR document specifies the details of the pediatric studies to be completed in order to receive the financial incentive (pediatric exclusivity).
- A sponsor may request the FDA to issue a WR by submission of a Proposed Pediatric Study Request (PPSR) <u>OR</u> FDA may issue a WR without a PPSR.
- PPSR should contain rationale for studies, detailed study designs and plans for formulation development



BPCA: Pediatric Exclusivity

- Applicants who fulfill the terms of a WR are eligible to receive pediatric exclusivity.
 - Additional 6 months of exclusivity
 - Exclusivity attaches to all existing marketing exclusivities and patents for the drug moiety.
 - Pediatric exclusivity does not require positive pediatric studies or granting a new indication.





Question: How many drugs have been approved for a pediatric cancer based on studies conducted under a Written Request?

- A: 2
- B: 17
- C: 0
- D: 9

BPCA facilitates Pediatric Oncology Drug Development



- Written Requests are the primary regulatory mechanism under BPCA for obtaining data for pediatric cancer drug approvals and product labeling.
- Since enactment of pediatric exclusivity legislation:
 - > 70 WRs issued by OHOP
 - > 9 drugs have been approved for pediatric cancer indications supported by studies conducted under a WR.
 - > 17 products have had pediatric safety/dosing information added to the label.
 - > 24 products have been granted exclusivity.
 - More than 30 products are currently being investigated under WR/BPCA.

Products with a pediatric oncology indication based on studies conducted under a WR



DRUG	CONDITION STUDIED UNDER WR	
Blinatumomab	ALL	
Clofarabine*	Refractory ALL	
Dasatinib	Ph+ CML in the chronic phase	
Everolimus	SEGA	
Imatinib	Ph+ ALL and Ph+ CML	
Ipilimumab	Unresectable or metastatic melanoma (> 12 years)	
Larotrectinib*	Metastatic or refractory solid tumors with NTRK gene fusion	
Nilotinib	Ph+ CML, relpased or refractory Ph+ ALL	
Tisagenlecleucel*	Relapsed or refractory ALL	

Products with pediatric information added to label based on a WR



DRUG	CONDITIONS STUDIED UNDER WR	
Bendamustine	Relapsed or refractory ALL and AML	
Bortezomib	Relapsed ALL	
Busulfan	Bone marrow transplant	
Cabazitaxel	High-grade glioma and DIPG	
Capecitabine	Newly diagnosed non-disseminated DIPG, HGG	
Carboplatin	Refractory or relapsed malignancy	
Docetaxel	Refractory or relapsed solid tumors	
Erlotinib	Refractory or relapsed ependymoma	
Fludarabine	Refractory acute leukemia	
Gemcitabine	Relapsed or refractory acute lymphoblastic leukemia	
Irinotecan	Refractory or relapsed solid tumors; newly diagnosed metastatic RMS	
Ixabepilone	Advanced or refractory solid tumors	
Oxaliplatin	Refractory or relapsed solid tumors	
Pemetrexed	Refractory or relapsed solid tumors	
Temsirolimus	Refractory or relapsed solid tumors; NBL, RMS, HGG	
Trabectedin	Histotypes of sarcoma, predominantly RMS, osteosarcoma, Ewing sarcoma, and non-RMS STS	
Vinorelbine	Leukemia or refractory or relapsed solid tumors	

Pediatric Labeling: (not so distant) Past and Present

Label Section	Asparaginase (Elspar)-2002	Tisagenlecleucel (Kymriah; CAR T-Cell)-2017
Indications and Usage	"in the therapy of patients with ALLuseful primarily in combination with other chemotherapeutic agents in the induction of remissions in pediatric patients"	"for the treatment of patients up to 25 years of age with B-cell precursor ALL that is refractory or in second or later relapse"

Pediatric Labeling: (not so distant) Past and Present



Label Section	Asparaginase (Elspar)-2002	Tisagenlecleucel (Kymriah; CAR T-Cell)-2017
Pediatric Use	"Asparaginase toxicity is reported to be greater in adults than in pediatric patients."	"The safety and efficacy of KYMRIAH have been established in pediatric patients with r/r B-cell ALL. Use of KYMRIAH is supported by a single-arm trial that included 52 pediatric patients with r/r B-cell precursor ALL in the following age groups: 33 children (age 3 years to less than 12 years) and 19 adolescents (age 12 years to less than 17 years). No differences in efficacy or safety were observed between the different age subgroups or in comparison to the young adults in the trial. "Reference to Sec 14.

Pediatric Labeling: (not so distant) Past and Present



Label Section	Asparaginase (Elspar)-2002	Tisagenlecleucel (Kymriah; CAR T-Cell)-2017
Clinical Studies	No description	The efficacy of KYMRIAH in pediatric and young adults with r/r B-cell precursor ALL was evaluated in an open-label, multicenter single-arm trial107 patients were screened, 88 were enrolled, 68 were treated, and 63 were evaluable for efficacy." The 63 evaluable patients included" age/gender/ethnicity/prior tx "The efficacy of KYMRIAH was established on the basis of" "Among the 63 infused patients", time to onset of remission, SCT rate Table summarizing efficacy results

Pediatric Subcommittee of the Oncologic Drugs Advisory Committee (ODAC)



- Industry sponsors obtain input from key academic and community opinion leaders regarding ongoing or potential pediatric development programs.
 - Gauge investigator interest in pediatric investigation of products in various stages of adult development
 - Provide feedback to industry on trial design, study population, pediatric regulations
 - Discuss drug candidates for a Written Request
- Interactive discussion of key and relevant topics
 - Immunotherapeutics, PRO instruments, FDARA legislation
- FDA invites sponsors of specific products early in development <u>AND</u> sponsors are encouraged to seek an invitation if there are questions regarding their pediatric development program.





Thanks!

