

Antiretroviral Drug Products Needed for Use Under PEPFAR¹

Single Drugs

Dolutegravir (pediatric formulations)

Two drug combinations with a protease inhibitor and ritonavir used as metabolic inhibitor (to be used in combination with other antiretroviral drugs)

darunavir/ritonavir

lopinavir/ritonavir (pediatric formulations)

Two and three-drug combinations (to be used in combination with other antiretroviral drugs)

abacavir + lamivudine (primarily for pediatric formulations)

tenofovir disoproxil fumarate (TDF) + emtricitabine (or lamivudine)

tenofovir alafenamide (TAF) + emtricitabine (or lamivudine)

Two and Three-drug combinations (complete regimen in treatment naïve)²

abacavir + lamivudine + dolutegravir

TDF (or TAF) + emtricitabine (or lamivudine) + efavirenz

TDF (or TAF) + emtricitabine (or lamivudine) + dolutegravir

Dolutegravir + lamivudine

Four Drug Combinations

Abacavir+lamivudine+lopinavir+ritonavir (pediatric formulations)

¹ This list is revised periodically to address current needs

² For any of the three drug combinations listed, any two components (in addition to those listed above) could be made into a fixed-combination that could be used in combination with other antiretroviral drugs. Some of these combinations may also be used in treatment-experienced patients.