

IMPORTANT PRESCRIBING INFORMATION

DATE: October 22, 2024

Subject: Temporary importation of EXTRANEAL (icodextrin) Peritoneal Dialysis Solution from Ireland for use in Automated Peritoneal Dialysis to address drug shortages

Dear Healthcare Professional,

Due to the current critical shortage of EXTRANEAL (icodextrin) Peritoneal Dialysis Solution in the United States (US) market, Baxter Healthcare Corporation (Baxter) is coordinating with the U.S. Food and Drug Administration (FDA) to temporarily import EXTRANEAL from Baxter's manufacturing facility in Castlebar, Ireland. FDA has not approved this product manufactured by Baxter's Castlebar, Ireland facility.

You may be provided with additional letters for other imported peritoneal dialysis solutions you receive. Please read each letter in its entirety because each letter may contain different, product-specific information.

Baxter has initiated temporary importation of EXTRANEAL (icodextrin) Peritoneal Dialysis Solution for use in Automated Peritoneal Dialysis (APD) therapy as described in the table below. This product is manufactured by Baxter's manufacturing facility in Castlebar, Ireland and is marketed in the United Kingdom and other countries within the European Union (EU). At this time, importation or distribution of EXTRANEAL (icodextrin) peritoneal dialysis solution in the United States by any entity other than Baxter or its authorized distributor(s) is considered a violation of the Federal Food, Drug, and Cosmetic Act and is subject to enforcement by the FDA.

Effective immediately, and during this temporary period, Baxter will offer the following imported products from Baxter's facility in Castlebar, Ireland:

Product Name and Description	APD Fill Volume	Product Code	Bags per Carton	NDC Code
EXTRANEAL (icodextrin) Solution for Peritoneal Dialysis	2000 mL	FPB4938RC	5 bags	0941-0719-01 (bag) 0941-0719-05 (carton)
EXTRANEAL (icodextrin) Solution for Peritoneal Dialysis	2500 mL	FPB4945R	4 bags	0941-0681-02 (bag) 0941-0681-08 (carton)

It is important to note the following:

- There are no clinically relevant differences in the EXTRANEAL drug composition between the
 European-manufactured and US-manufactured APD product (see Table 1). As such, clinical practice
 for usage, administration, and dosage for Extraneal with 7.5% icodextrin (manufactured in EU)
 products is the same as with the Extraneal with 7.5% icodextrin (manufactured in U.S.). Please refer
 to the FDA-approved EXTRANEAL (icodextrin) Peritoneal Dialysis Solution Prescribing Information
 for reference.
- Calcium and Magnesium electrolyte concentrations are identical in EXTRANEAL manufactured in the EU and US but appear different as they are expressed in mmol/L (EU) and in mEq/L (U.S.).
- The Luer-lock connector on the Ireland imported product functions the same as and is fully compatible with peritoneal dialysis sets marketed in the United States. However, the U.S. product has a purple pull ring



covering the luer to identify the solution, while the Ireland imported product has a blue protective tip protector. The frangible is green in the Ireland imported product but blue in U.S. product. Users of the imported product should check the product label to ensure that they are using the correct APD solution. See Table 1 for more details of product differences.

EXTRANEAL (icodextrin) solution for peritoneal dialysis imported carton labeling includes barcodes; however, the
barcodes may not register accurately in the US scanning systems. There are no barcodes on the solution
containers of the Ireland imported product. Alternative procedures should be followed to assure that the correct
drug product is being used in all systems and processes and administered to individual patients. For example,
institutions should consider manually inputting the product into their systems and confirm that barcode systems do
not provide incorrect information when the product is scanned.

Before prescribing, healthcare providers should be aware of some key differences in the container packaging and labeling between the EXTRANEAL (icodextrin) Solution for Peritoneal Dialysis products (manufactured in EU) and EXTRANEAL (icodextrin) Peritoneal Dialysis Solution (manufactured in U.S.).

Key differences are highlighted in the following Product Comparison Tables:

- Table 1: Key differences of EXTRANEAL for APD therapy
- Table 2: Label images of EXTRANEAL for 2000 mL and 2500 mL APD product presentations

Reporting Adverse Events

To report adverse events associated with the imported product, please call Baxter at 1-866-888-2472, or fax: 1-800-759-1801. Adverse events or quality problems experienced with the use of the imported product may also be reported to the FDA's MedWatch Adverse Event Reporting program either online, by regular mail or by fax:

- Complete and submit the report Online: www.fda.gov/medwatch/report.htm
- Regular mail or Fax: Download form www.fda.gov/MedWatch/getforms.htm or call 1-800-332-1088 to request a reporting form, then complete and return to the address on the pre-addressed form or submit by fax to 1-800-FDA-0178 (1-800-332-0178).

To report product quality issues, please report to:

Baxter - Product Feedback Portal (https://productfeedback.baxter.com/).

Please refer to the FDA approved prescribing information for EXTRANEAL (icodextrin) Peritoneal Dialysis Solution at DailyMed (nih.gov).

If you have any questions about the information contained in this letter or the use of imported EXTRANEAL (icodextrin) Solution for Peritoneal Dialysis, please contact Baxter's Medical Information Service at 1-888-736-2543.

To place an order, please contact Baxter's Center for Home Care Services by calling 1-800-284-4060.



Sincerely,

Electronically signed by: Geovana Basso Reason: Ok. CDT

Electronically signed by: Geovana Basso Reason: Ok. CDT

Geovana Basso, M.D.
Director of Americas Medical Affairs
Baxter Healthcare Corporation
One Baxter Parkway
Deerfield, Illinois 60015

Baxter, Ambu-Flex and EXTRANEAL are registered trademarks of Baxter International Inc.

Attachments:

Product Comparison Tables 1 and 2



Table 1. Key differences of EXTRANEAL for APD therapy

	Imported Product from Castlebar, Ireland	U.S. FDA Approved Product
Product name	EXTRANEAL (icodextrin) Solution for Peritoneal Dialysis	EXTRANEAL (icodextrin) Peritoneal Dialysis Solution
Labeled Fill Volume	2000 mL 2500 mL	2000 mL 2500 mL
Container Type	PVC Container	Ambu-Flex Container (PVC)
Bags per carton	2000 mL: 5 bags 2500 mL:4 bags	2000 mL: 6 bags 2500 mL: 5 bags
Indications	Extraneal is recommended as a once daily replacement for a single glucose exchange as part of a continuous ambulatory peritoneal dialysis (CAPD) or automated peritoneal dialysis (APD) regimen for the treatment of chronic renal failure, particularly for patients who have lost ultrafiltration on glucose solutions, because it can extend time on CAPD therapy in such patients.	EXTRANEAL (icodextrin) is indicated for a single daily exchange for the long (8- to 16- hour) dwell during continuous ambulatory peritoneal dialysis (CAPD) or automated peritoneal dialysis (APD) for the management of kidney failure in patients requiring long-term kidney replacement therapy. EXTRANEAL is also indicated to improve (compared to 4.25% dextrose) long-dwell ultrafiltration and clearance of creatinine and urea nitrogen in patients with high average or greater transport characteristics, as defined using the peritoneal equilibration test (PET)
Active Ingredients	75 g/L Icodextrin (7.5 g/100 mL) 5.4 g/L Sodium Chloride (540 mg/100 mL) 4.5 g/L Sodium Lactate (450 mg/100 mL) 0.257g/L Calcium Chloride (25.7 mg/100 mL) 0.051g/L Magnesium Chloride (5.10 mg/100 mL)	7.5 g/100 mL Icodextrin 535 mg/100 mL Sodium Chloride, USP* 448 mg/100 mL Sodium Lactate* 25.7 mg/100 mL Calcium Chloride, USP* 5.08 mg/100 mL Magnesium Chloride, USP* * considered excipients in US drug registration
Electrolyte Content per Liter	Sodium 133 mmol/L (equivalent to 133 mEq/L) Calcium 1.75 mmol/L (equivalent to 3.5 mEq/L) Magnesium 0.25 mmol/L (equivalent to 0.5 mEq/L) Chloride 96 mmol/L (equivalent to 96 mEq/L) Lactate 40 mmol/L (equivalent to 40 mEq/L)	Sodium 132 mEq/L Calcium 3.5 mEq/L Magnesium 0.5 mEq/L Chloride 96 mEq/L Lactate 40 mEq/L
рH	pH 5.0 – 6.0 HCl / NaOH may have been used to adjust pH	pH 5.0 – 6.0 HCl / NaOH may have been used to adjust pH
Additional Information	Osmolarity 284 mOsm/L	Osmolarity (Calc) 282 – 286 mOsmol/L
Storage Conditions	Do not store below 4°C	Store at 20–25°C (68–77°F). Excursions permitted to 15–30°C (59–86°F) [See USP Controlled Room Temperature]. Protect from freezing.
Expiration Dating	24 months	18 months




	Imported Product from Castlebar, Ireland	U.S. FDA Approved Product
Container Closure System	Company Test Company C	Service of the servic
Container Closure	T)	
Container Closure System Differences	 One green frangible at luer-lock connector Blue protective tip protector 	 One blue frangible at luer-lock connector Purple pull ring cap closure



Table 2. Comparison of EXTRANEAL (icodextrin) PD Solution Container Labels

A comparison of labels is provided below. Note, both Baxter and Vantive-branded labels are presented, which represent all imported lot labeling scenarios during this transition.





Imported Product from Ca	astlebar, Ir	eland		US FDA approved Product
Lot			EIP	
B4938R			2000 ml	
Vantive				
	I Dielecie			
EXTRANEAL Solution for Peritonea	II DIAIYSIS	; 		
Formula 1000 ml			mmol/I	
Icodextrin	75 g		133	
Sodium Chloride	5,4 g		1,75	
Sodium (S)-lactate	4,5 g	Mg ⁺⁺	0,25	
Calcium Chloride 2H ₂ 0	0,257 g	CI-	96	
Magnesium Chloride 6H ₂ 0	0,051 g	С ₃ H ₅ О ₃	40	
Water for Injections		Osmolarity	284 m0sm/l	
Hydrochloric acid or sodium hydroxide (for pH adjustment)		pН	5,0 - 6,0	
Not for intravenous use Use as directed by a physician Read the package leaflet before use Knep out of the sight and reach of children Nonsyrogenic Do not store below 4°C Do not use unless solution is clear and centainer undamaged For single use only Once removed from the overpouch use immediately Discard unused solution Vantire Limited Wavertree Boulevard Liverpool, I7 oPPE United Kingdom Pt. 58711/0005				
POM			C8-35-05-248	
VANTIVE VE	RSION			



Calcium Chloride 2H₂O 0,257 g cr 96 Magnesium Chloride 6H₂O 40 0,051 g C3H5O3 Water for Injections 284 m0sm/l **Osmolarity** Hydrochloric acid or sodium hydroxide (for pH adjustment) рΗ 5,0-6,0

For intraperitoneal use Not for intravenous use Use as directed by a physician Read the package leaflet before use Keep out of the sight and reach of children

Nonpyrogenic Do not store below 4°C

Do not use unless solution is clear and container undamaged

For single use only
Once removed from the overpouch use immediately

Discard unused solution

Baxter Healthcare Ltd Caxton Way, Thetford, Norfolk, IP24 3SE, United Kingdom

Manufactured by Baxter Healthcare S A IRL-Castlebar CB-35-04-606

PL 00116/0266 PA 2299/017/001 MA 1277/01101 POM

BAXTER VERSION

IE and MT: Baxter Holding B.V.,

Kobaltweg 49, 3542CE, Utrecht, Netherlands

US FDA approved Product

L5B4976 NDC 0941-0679-05

0

2500⊾ (APPROX 90 mL EXCESS)

Baxter

Extraneal (icodextrin) Peritoneal Dialysis Solution



EACH 100 mL CONTAINS 7.5 g ICODEXTRIN 535 mg SODIUM CHLORIDE USP 448 mg SODIUM LACTATE 25.7 mg CALCIUM CHLORIDE USP 5.08 mg MAGNESIUM CHLORIDE USP WATER FOR INJECTION USP

mEq/L SODIUM 132 CALCIUM 3.5 MAGNESIUM 0.5 CHLORIDE 96 LACTATE 40 PH 5.0 - 6.0 PH MAY HAVE BEEN ADJUSTED WITH HYDROCHLORIC ACID OR SODIUM HYDROXIDE

EXTRANEAL SOLUTION CONTAINS NO BACTERIOSTATIC OR ANTIMICROBIAL AGENTS OSMOLARITY (CALC) 292 - 296 mOsmol/L

STERILE NONPYROGENIC

POTASSIUM CHLORIDE TO BE ADDED ONLY UNDER THE DIRECTION OF A PHYSICIAN

SEE PACKAGE INSERT FOR DOSAGE INFORMATION

USE AS DIRECTED BY PHYSICIAN

FOR INTRAPERITONEAL ADMINISTRATION ONLY

CAUTIONS SQUEEZE AND INSPECT INNER BAG THAT MAINTAINS PRODUCT STERILITY DISCARD IF LEAKS ARE

DO NOT USE UNLESS SOLUTION IS CLEAR

DISCARD UNUSED PORTION

BX ONLY

STORE IN MOISTURE BARRIER OVERPOUCH IN CARTON UNTIL READY TO USE

STORE AT 20-25°C (68-77°F) EXCURSIONS PERMITTED TO 15-30°C (59-86°F) [SEE USP CONTROLLED ROOM TEMPERATURE] PROTECT FROM FREEZING

PL 146 PLASTIC

Ambu-Flex II CONTAINER

BAXTER EXTRANEAL AMBU-FLEX II AND PL 146 ARE TRADEMARKS OF BAXTER INTERNATIONAL INC

BAXTER HEALTHCARE CORPORATION

DEERFIELD IL 60015 USA

MADE IN USA

US PAT NOS 4761237 4886789 6077836 6248726 B1



Imported Product from C	astlebar, lı	reland		US FDA approved Product
Lot B4945R			2500 ml	
Vantive				
EXTRANEAL Solution for Peritonea	l Dialysis			
Formula 1000 ml			mmol/l	
Icodextrin	75 g	Na+	133	
Sodium Chloride	5,4 g		1,75	
Sodium (S)-lactate	4,5 g	Mg++	0,25	
Calcium Chloride 2H ₂ 0	0,257 g		96	
Magnesium Chloride 6H ₂ O	0,051 g		40	
Water for Injections		Osmolarity	284 m0sm/l	
Hydrochloric acid or sodium hydroxide (for pH adjustment)		pH	5,0-6,0	
For intraperitoneal use Not for intraperitoneal use Not for intravenous use Use as directed by a physician Read the package leaflet before use Keep out of the sight and reach of children Nonpyrogenic Do not shore below 4°C Do not use unless solution is clear and container undamaged For single use only Once removed from the overpouch use immediately Discard unused solution Vantive Limited Wavertree Boulevard Liverpoot, I.7 9PE United Kingdom PL 5 871 1 / 000 5				
POM			00 05 05 C10	
VANTIVE VE	RSION		CB-35-05-249	