

TTT # 2022-2468

NDA 020687

ANDA 091178

**Mifepristone U.S. Post-Marketing Adverse Events Summary through 12/31/2024**

The following information is from United States (U.S.) post-marketing reports received by FDA of adverse events that occurred among patients who had taken mifepristone for medical termination of pregnancy. Because FDA has eliminated duplicate reports, and in some cases, reclassified the adverse event terms for individual cases after reviewing the narrative details, the numbers provided here may differ from the numbers of the reports that may be obtained through Freedom of Information Act requests. These events cannot with certainty be causally attributed to mifepristone because of information gaps about patient health status, clinical management of the patient, concurrent drug use, and other possible medical or surgical treatments and conditions. The estimated number of women who have used mifepristone in the U.S. for medical termination of pregnancy through the end of December 2024 is approximately 7.5 million women.

For informational purposes, fatal foreign cases that were reported after U.S. approval of mifepristone for medical termination of pregnancy are also included in a footnote in Table 1.

Date range of cumulative reports	09/28/00 <sup>†</sup> - 12/31/24
Died <sup>‡</sup>	36
*Ectopic pregnancies <sup>  </sup>	97
<p><sup>†</sup> U.S. approval date</p> <p><sup>‡</sup> The fatal cases are included regardless of causal attribution to mifepristone. Deaths were associated with sepsis in 13 of the 36 reported fatalities (9 cases tested positive for <i>Clostridium sordellii</i>, 1 case tested positive for <i>Clostridium perfringens</i>, 2 cases had negative blood cultures, and 1 case did not have blood culture data). Ten of the 13 fatal sepsis cases reported vaginal misoprostol use; 2 cases reported buccal misoprostol use; 1 case did not report the route of misoprostol use. Twenty-two of the 23 remaining U.S. deaths involved 2 cases of homicide, 2 cases of combined drug intoxication/overdose, 2 cases of ruptured ectopic pregnancy, 2 cases of drug intoxication, 2 cases of suicide, and 1 case each of the following: substance abuse/drug overdose; methadone overdose; suspected homicide; delayed onset toxic shock-like syndrome; hemorrhage; bilateral pulmonary thromboemboli; unintentional overdose resulting in liver failure; probable anaphylactic medication reaction; septic shock due to necrotizing fasciitis; sepsis with multiple complications possibly secondary to toxic shock syndrome 82 days after mifepristone; sudden death of undetermined etiology despite performance of an autopsy; and a case of natural death due to severe pulmonary emphysema. In the 23<sup>rd</sup> case, the cause of death could not be established despite performance of an autopsy; tissue samples were negative for <i>Clostridium sordellii</i>. There were 13 additional reported deaths in women in foreign countries who used mifepristone for medical termination of pregnancy. These fatal cases were associated with the following: sepsis (<i>Clostridium sordellii</i> identified in tissue samples) in a foreign clinical trial; sepsis (Group A <i>Streptococcus pyogenes</i>); a ruptured gastric ulcer; severe hemorrhage; severe hemorrhage and possible sepsis; "multivisceral failure;" thrombotic thrombocytopenic purpura leading to intracranial hemorrhage; toxic shock syndrome (<i>Clostridium sordellii</i> was identified through uterine biopsy cultures); sepsis (<i>Enterococcus faecalis</i> and <i>Escherichia coli</i> were identified in blood culture); asthma attack with cardiac arrest; thromboembolism; respiratory decompensation with secondary pulmonary infection 30 days after mifepristone in a patient on the lung transplant list with diabetes, a jejunostomy feeding tube, and severe cystic fibrosis; and a case of <i>Clostridium septicum</i> sepsis (from a published literature report).</p> <p>* The majority of these women are included in the hospitalized category in Table 2.</p> <p><sup>  </sup> Administration of mifepristone and misoprostol is contraindicated in patients with confirmed or suspected ectopic pregnancy (a pregnancy outside the uterus).</p>	

<b>Table 2. Post-Marketing Adverse Events in U.S. Women Who Used Mifepristone for Medical Termination of Pregnancy</b>		
Date ranges of reports received	09/28/00 <sup>†</sup> - 10/31/12	11/01/12 - 12/31/24 <sup>‡</sup>
Cases with any adverse event	2740	1512
Hospitalized, excluding deaths	768	288
*Experienced blood loss requiring transfusions <sup>§</sup>	416	190
Infections <sup>  </sup> (*Severe infections <sup>¶</sup> )	308 (57)	114 (22)

<sup>†</sup> U.S. approval date

<sup>‡</sup> FDA implemented the FDA Adverse Event Reporting System (FAERS) on September 10, 2012, and migrated all the data from the previous reporting system (AERS) to FAERS. Differences may exist when comparing case counts in AERS and FAERS. FDA validated and recoded product information as the AERS reports were migrated to FAERS. As a result of this change, it is not recommended to calculate a cumulative number when reviewing the data provided in Table 2.

\* The majority of these women are included in the hospitalized category in Table 2.

<sup>§</sup> As stated in the approved labeling for Mifeprex (mifepristone) and its approved generic version, bleeding or spotting can be expected for an average of 9-16 days, and may last for up to 30 days. Excessive vaginal bleeding usually requires treatment by uterotonics, vasoconstrictor drugs, curettage, administration of saline infusions, and/or blood transfusions.

<sup>||</sup> This category includes endometritis (inflammation resulting from an infection involving the lining of the womb), pelvic inflammatory disease (involving the nearby reproductive organs such as the fallopian tubes or ovaries), and pelvic infections with sepsis (a serious systemic infection that has spread beyond the reproductive organs). Not included are women with reported sexually transmitted infections such as chlamydia and gonorrhea, cystitis, and toxic shock syndrome not associated with a pelvic infection.

<sup>¶</sup> This subset of infections includes cases that were determined to be severe based on medical review of the available case details. Severe infections generally result in death or hospitalization for at least 2-3 days, require intravenous antibiotics for at least 24 hours and total antibiotic usage for at least 3 days, or have other physical or clinical findings, laboratory data, or surgery that suggest a severe infection.