

BAM: Salmonella

December 2007 Edition*

Bacteriological Analytical Manual Chapter 5 *Salmonella*

Authors

Chapter Contents

- [Introduction](#)
- [Equipment and Materials](#)
- [Media and Reagents](#)
- [Preparation of foods for isolation of *Salmonella*](#)
- [Isolation of *Salmonella*](#)
- [Identification of *Salmonella*](#)
- [Rapid methods \[Appendix 1, see below\]](#)
- [References](#)

Introduction

Several changes are being introduced in this edition of BAM (8th Edition). The first change involves the expanded use of [Rappaport-Vassiliadis \(RV\) medium](#) for foods with both high and low levels of competitive microflora. In the previous edition, RV medium was recommended only for the analysis of shrimp. Based on the completion of AOAC precollaborative (5, 6) and collaborative (7, 8) studies, RV medium is now being recommended for the analysis of high microbial and low microbial load foods. RV medium replaces selenite cystine (SC) broth for the analysis of all foods, except guar gum. In addition, RV medium replaces lauryl tryptose broth for use with dry active yeast. [Tetrathionate \(TT\)](#) broth continues to be used as the second selective enrichment broth. However, TT broth is to be incubated at 43°C for the analysis of high microbial load foods and at 35°C for the analysis of low microbial load foods, including guar gum.

The second change involves the option of refrigerating incubated preenrichments and selective enrichments of low-moisture foods for up to 72 h. With this option, sample analyses can be initiated as late as Wednesday or Thursday without weekend work being involved.

The third change involves reducing the period of incubation of the [lysine iron agar \(LIA\)](#) slants. In the former edition (BAM-7), [triple sugar iron agar \(TSI\)](#) and LIA slants were incubated at 35°C for 24 ± 2 h and 48 ± 2 h, respectively. Unpublished data have demonstrated that the 48 h reading of LIA slants is without diagnostic value. Of 193 LIA slants examined, all gave definitive results within 24 ± 2 h of incubation. No significant changes altered the final test result when the slants were incubated an additional 24 h. Thus, both the TSI and LIA slants are now incubated for 24 ± 2 h.

The fourth change involves the procedure for surface disinfection of shell eggs. In the previous edition (BAM-7), egg shells were surface-disinfected by soaking in 0.1% mercuric chloride solution for 1 h followed by soaking in 70% ethanol for 30 min. Mercuric chloride is classified as a hazardous waste, and is expensive to dispose of according to Environmental Protection Agency guidelines. In this edition (BAM-8) egg shells are now surface-disinfected by soaking for at least 10 sec in a 3:1 solution consisting of 3 parts of 70% alcohol (ethyl or isopropyl) to 1 part of iodine/potassium iodide solution.

The fifth change involves the sample preparation of eggs. Egg contents (yolk and albumen) are thoroughly mixed before analysis. After mixing the egg contents, 25 g (ml) are added to 225 ml trypticase (tryptic) soy broth supplemented with ferrous sulfate.

A method for the analysis of guar gum has been included. When guar gum is preenriched at a 1:9 sample/broth ratio, a highly viscous, nonpipettable mixture results. Addition of the enzyme cellulase to the preenrichment medium, however, results in a readily pipettable mixture.

A method for orange juice (pasteurized and unpasteurized) has been included due to recent orange juice-related outbreaks.

The directions for picking colonies from the selective plating agars have been made more explicit to reflect the intent of the method. In the absence of typical or suspect colonies on the selective plating agars, it is recommended that atypical colonies be picked to TSI and LIA slants. This recommendation is based on the fact that up to 4% of all *Salmonella* cultures isolated by FDA analysts from certain foods, especially seafoods, during the past several years have been atypical.

Finally, since the publication of BAM-7, a 6-way comparison was conducted of the relative effectiveness of the three selective plating agars recommended in the BAM ([bismuth sulfite](#), [Hektoen enteric](#), and [xylose lysine desoxycholate agars](#)) and three relatively new agars (EF-18, xylose lysine Tergitol 4, and Rambach agars). Our results (9) indicated no advantage in replacing any of the BAM-recommended agars with one or more of the newer agars. Thus, the combination of selective plating agars recommended in BAM-7 remains unchanged.

Return to Chapter Contents

A. Equipment and materials

1. Blender and sterile blender jars (**see** Chapter 1)
2. Sterile, 16 oz (500 ml) wide-mouth, screw-cap jars, sterile 500 ml Erlenmeyer flasks, sterile 250 ml beakers, sterile glass or paper funnels of appropriate size, and, optionally, containers of appropriate capacity to accommodate composited samples
3. Sterile, bent glass or plastic spreader rods
4. Balance, with weights; 2000 g capacity, sensitivity of 0.1 g
5. Balance, with weights; 120 g capacity, sensitivity of 5 mg
6. Incubator, $35 \pm 2^\circ\text{C}$
7. Refrigerated incubator or laboratory refrigerator, $4 \pm 2^\circ\text{C}$
8. Water bath, $49 \pm 1^\circ\text{C}$
9. Water bath, circulating, thermostatically-controlled, $43 \pm 0.2^\circ\text{C}$
10. Water bath, circulating, thermostatically-controlled, $42 \pm 0.2^\circ\text{C}$
11. Sterile spoons or other appropriate instruments for transferring food samples
12. Sterile culture dishes, 15 x 100 mm, glass or plastic
13. Sterile pipets, 1 ml, with 0.01 ml graduations; 5 and 10 ml, with 0.1 ml graduations
14. Inoculating needle and inoculating loop (about 3 mm id or 10 5l), nichrome, platinum-iridium, chromel wire, or sterile plastic
15. Sterile test or culture tubes, 16 x 150 mm and 20 x 150 mm; serological tubes, 10 x 75 mm or 13 x 100 mm
16. Test or culture tube racks
17. Vortex mixer
18. Sterile shears, large scissors, scalpel, and forceps
19. Lamp (for observing serological reactions)
20. Fisher or Bunsen burner
21. pH test paper (pH range 6-8) with maximum graduations of 0.4 pH units per color change
22. pH meter
23. Plastic bags, 28 x 37 cm, sterile, with resealable tape. (Items 23-24 are needed in the analysis of frog legs and rabbit carcasses.)

24. Plastic beakers, 4 liter, autoclavable, for holding plastic bag during shaking and incubation.
25. Sponges, non-bactericidal (Nasco cat # B01299WA), or equivalent.
26. Swabs, non-bactericidal, cotton-tipped.

Return to Chapter Contents

B. [Media and reagents](#)

For preparation of media and reagents, refer to Methods 967.25-967.28 in *Official Methods of Analysis* (1).

1. Lactose broth ([M74](#))
2. Nonfat dry milk (reconstituted) ([M111](#))
3. Selenite cystine (SC) broth ([M134](#))
4. Tetrathionate (TT) broth ([M145](#))
5. Rappaport-Vassiliadis (RV) medium ([M132](#)). NOTE: RV medium must be made from its individual ingredients. Commercial formulations are not acceptable.
6. Xylose lysine desoxycholate (XLD) agar ([M179](#))
7. Hektoen enteric (HE) agar ([M61](#))
8. Bismuth sulfite (BS) agar ([M19](#))
9. Triple sugar iron agar (TSI) ([M149](#))
10. Tryptone (tryptophane) broth ([M164](#))
11. Trypticase (tryptic) soy broth ([M154](#))
12. Trypticase soy broth with ferrous sulfate ([M186](#))
13. Trypticase soy-tryptose broth ([M160](#))
14. MR-VP broth ([M104](#))
15. Simmons citrate agar ([M138](#))
16. Urea broth ([M171](#))
17. Urea broth (rapid) ([M172](#))
18. Malonate broth ([M92](#))
19. Lysine iron agar (LIA) (Edwards and Fife) ([M89](#))
20. Lysine decarboxylase broth ([M87](#))
21. Motility test medium (semisolid) ([M103](#))
22. Potassium cyanide (KCN) broth ([M126](#))
23. Phenol red carbohydrate broth ([M121](#))
24. Purple carbohydrate broth ([M130](#))
25. MacConkey agar ([M91](#))
26. Nutrient broth ([M114](#))
27. Brain heart infusion (BHI) broth ([M24](#))
28. Papain solution, 5% ([M56a](#))
29. Cellulase solution, 1% ([M187](#))
30. Tryptose blood agar base ([M166](#))
31. Universal preenrichment broth ([M188](#))
32. Universal preenrichment broth (without ferric ammonium citrate) ([M188a](#))
33. Buffered peptone water ([M192](#))
34. Dey-Engley broth ([M193](#))
35. Potassium sulfite powder, anhydrous
36. Chlorine solution, 200 ppm, containing 0.1% sodium dodecyl sulfate ([R12a](#))
37. Ethanol, 70% ([R23](#))
38. Kovacs' reagent ([R38](#))
39. Voges-Proskauer (VP) test reagents ([R89](#))
40. Creatine phosphate crystals
41. Potassium hydroxide solution, 40% ([R65](#))
42. 1 N Sodium hydroxide solution ([R73](#))
43. 1 N Hydrochloric acid ([R36](#))
44. Brilliant green dye solution, 1% ([R8](#))

45. Bromcresol purple dye solution, 0.2% ([R9](#))
46. Methyl red indicator ([R44](#))
47. Sterile distilled water
48. Tergitol Anionic 7 ([R78](#))
49. Triton X-100 ([R86](#))
50. Physiological saline solution, 0.85% (sterile) ([R63](#))
51. Formalinized physiological saline solution ([R27](#))
52. *Salmonella* polyvalent somatic (O) antiserum
53. *Salmonella* polyvalent flagellar (H) antiserum
54. *Salmonella* somatic group (O) antisera: A, B, C₁, C₂, C₃, D₁, D₂, E₁, E₂, E₃, E₄, F, G, H, I, Vi, and other groups, as appropriate
55. *Salmonella* Spicer-Edwards flagellar (H) antisera

[Return to Chapter Contents](#)

C. [Preparation of foods for isolation of *Salmonella*](#)

The following methods are based on the analysis of a 25 g analytical unit at a 1:9 sample/broth ratio. Depending on the extent of compositing, add enough broth to maintain this 1:9 ratio unless otherwise indicated. For samples not analyzed on an exact weight basis, e.g., frog legs, refer to the specific method for instructions.

1. **Dried egg yolk, dried egg whites, dried whole eggs, liquid milk (skim milk, 2% fat milk, whole, and buttermilk), and prepared powdered mixes (cake, cookie, doughnut, biscuit, and bread), infant formula, and oral or tube feedings containing egg.**

Preferably, do not thaw frozen samples before analysis. If frozen sample must be tempered to obtain analytical portion, thaw suitable portion as rapidly as possible to minimize increase in number of competing organisms or to reduce potential of injuring *Salmonella* organisms. Thaw below 45°C for 15 min with continuous agitation in thermostatically controlled water bath or thaw within 18 h at 2-5°C. Aseptically weigh 25 g sample into sterile, wide-mouth, screw-cap jar (500 ml) or other appropriate container. For nonpowdered samples, add 225 ml sterile [lactose broth](#). If product is powdered, add about 15 ml sterile lactose broth and stir with sterile glass rod, spoon, or tongue depressor to smooth suspension. Add 3 additional portions of lactose broth, 10, 10, and 190 ml, for total of 225 ml. Stir thoroughly until sample is suspended without lumps. Cap jar securely and let stand 60 ± 5 min at room temperature. Mix well by swirling and determine pH with test paper. Adjust pH, if necessary, to 6.8 ± 0.2 with sterile 1 N NaOH or 1 N HCl. Cap jar securely and mix well before determining final pH. Loosen jar cap about 1/4 turn and incubate 24 ± 2 h at 35°C. Continue as in D, 1-11, below.

2. **Eggs**
 - a. **Shell eggs.** Remove any adherent material from the shell surface. Disinfect eggs with 3:1 solution consisting of 3 parts of 70% alcohol (ethyl or isopropyl) to 1 part iodine/potassium iodide solution. Prepare 70% alcohol solution either by diluting 700 ml 100% alcohol with sterile distilled water for a final volume of 1,000 ml or by diluting 700 ml 95% alcohol with sterile distilled water for a final volume of 950 ml. Prepare iodine/potassium iodide solution by dissolving 100 g potassium iodide in 200-300 ml sterile distilled water. Add 50 g iodine and heat gently with constant mixing until the iodine is dissolved. Dilute the iodine/potassium iodide solution to 1,000 ml with sterile distilled water. Store iodine/potassium iodide solution in amber glass-stoppered bottle in the dark. Prepare the disinfection solution by adding 250 ml iodine/potassium iodide solution to 750 ml 70% alcohol solution and mix well. Submerge eggs in disinfection solution for at least 10 seconds. Remove eggs and allow to air dry. Eggs with chipped, cracked, or

- broken shells are not included in the sample. Each sample shall consist of twenty (20) eggs cracked aseptically into a Whirl-Pak bag, for a total of fifty (50) samples per poultry house. Eggs are cracked aseptically by gloved hands, with a change of gloves between samples. Mix samples thoroughly by gloved hands, with a change of gloves between samples. Mix samples thoroughly by hand until yolks are completely mixed with the albumen. Samples are held at room temperature (20-24°C) for 96 ± 2 h. After 96 ± 2 h, remove 25 ml portion from each sample of pooled eggs, and preenrich 25 ml test portion in 225 ml sterile [trypticase soy broth \(TSB\) supplemented with ferrous sulfate](#) (35 mg ferrous sulfate added to 1000 ml TSB) and mix well by swirling. Let stand 60 ± 5 min at room temperature. Mix well by swirling and determine pH with test paper. Adjust pH, if necessary, to 6.8 ± 0.2. Incubate 24 ± 2 h at 35°C. Continue as in D, 1-11, below.
- b. **Liquid whole eggs (homogenized).** Combine fifteen (15) 25 ml test portions into a 375 ml composite contained in a 6-liter Erlenmeyer flask. Composites are held at room temperature (20-24°C) for 96 ± 2 h. After 96 ± 2 h, add 3,375 ml sterile [TSB supplemented with ferrous sulfate](#), as described above, and mix well by swirling. Let stand 60 ± 5 min at room temperature. Mix well by swirling and determine pH with test paper. Adjust pH, if necessary, to 6.8 ± 0.2. Incubate 24 ± 2 h at 35°C. Continue as in D, 1-11, below.
 - c. **Hard-boiled eggs (chicken, duck, and others).** If the egg shells are still intact, disinfect the shells as described above and aseptically separate the shells from the eggs. Pulverize the eggs (egg yolk solids and egg white solids) aseptically and weigh 25 g into a sterile 500 ml Erlenmeyer flask or other appropriate container. Add 225 ml [TSB](#) (without ferrous sulfate) and mix well by swirling. Continue as described above.
3. **Nonfat dry milk**
 - a. **Instant.** Aseptically weigh 25 g sample into sterile beaker (250 ml) or other appropriate container. Using sterile glass or paper funnel (made with tape to withstand autoclaving), pour 25 g analytical unit gently and slowly over surface of 225 ml brilliant green water contained in sterile 500 ml Erlenmeyer flask or other appropriate container. Alternatively, 25 g analytical units may be composited and poured over the surface of proportionately larger volumes of brilliant green water. Prepare brilliant green water by adding 2 ml [1% brilliant green dye solution](#) per 1000 ml sterile distilled water. Let container stand undisturbed for 60 ± 5 min. Incubate loosely capped container, without mixing or pH adjustment, for 24 ± 2 h at 35°C. Continue as in D, 1-11, below.
 - b. **Non-Instant.** Examine as described for instant nonfat dry milk, except that the 25 g analytical units may not be composited.
 4. **Dry whole milk.** Examine as described for instant nonfat dry milk, except that the 25 g analytical units may not be composited.
 5. **Casein**
 - a. **Lactic casein.** Aseptically weigh 25 g sample into sterile beaker (250 ml) or other appropriate container. Using sterile glass or paper funnel (made with tape to withstand autoclaving), pour 25 g analytical unit gently and slowly over the surface of 225 ml Universal Preenrichment broth contained in sterile 500 ml Erlenmeyer flask or other appropriate container. Analytical units (25 g) may be composited. Let container stand undisturbed 60 ± 5 min. Incubate loosely capped container, without mixing or pH adjustment, for 24 ± 2 h at 35°C. Continue as in D, 1-11, below.
 - b. **Rennet casein.** Aseptically weigh 25 g sample into sterile beaker (250 ml) or other appropriate container. Using sterile glass or paper funnel (made with tape to withstand autoclaving), pour 25 g analytical unit gently and slowly over the surface of 225 ml lactose broth contained in sterile 500 ml Erlenmeyer flask or other appropriate container. Analytical units (25 g) may be composited. Let container stand undisturbed 60 ± 5 min. Incubate loosely capped container, without mixing or pH adjustment, for 24 ± 2 h at 35°C. Continue as in D, 1-11, below.

- c. **Sodium caseinate.** Aseptically weigh 25 g sample into sterile, wide-mouth, screw-cap jar (500 ml) or other appropriate container. Add 225 ml sterile lactose broth and mix well. Analytical units may be composited. Let stand 60 min at room temperature with jar securely capped. Mix well by swirling and determine pH with test paper. Adjust pH, if necessary, to 6.8 ± 0.2 . Loosen jar about 1/4 turn and incubate 24 ± 2 h at 35°C . Continue as in D, 1-11, below.
- 6. **Soy flour.** Examine as described for rennet casein, except 25 g analytical units (25 g) may not be composited.
- 7. **Egg-containing products (noodles, egg rolls, macaroni, spaghetti), cheese, dough, prepared salads (ham, egg, chicken, tuna, turkey), fresh, frozen, or dried fruits and vegetables, nut meats, crustaceans (shrimp, crab, crayfish, langostinos, lobster), and fish.** Preferably, do not thaw frozen samples before analysis. If frozen sample must be tempered to obtain analytical portion, thaw below 45°C for <15 min with continuous agitation in thermostatically controlled water bath or thaw within 18 h at $2\text{-}5^{\circ}\text{C}$.

Aseptically weigh 25 g sample into sterile blending container. Add 225 ml sterile [lactose broth](#) and blend 2 min. Aseptically transfer homogenized mixture to sterile, wide-mouth, screw-cap jar (500 ml) or other appropriate container and let stand 60 ± 5 min at room temperature with jar securely capped. Mix well by swirling and determine pH with test paper. Adjust pH, if necessary, to 6.8 ± 0.2 . Mix well and loosen jar cap about 1/4 turn. Incubate 24 ± 2 h at 35°C . Continue as in D, 1-11, below.

- 8. **Dried yeast (active and inactive yeast).** Aseptically weigh 25 g sample into sterile, wide-mouth, screw-cap jar (500 ml) or other appropriate container. Add 225 ml sterile [trypticase soy broth](#). Mix well to form smooth suspension. Let stand 60 ± 5 min at room temperature with jar securely capped. Mix well by swirling and determine pH with test paper. Adjust pH, if necessary, to 6.8 ± 0.2 , mixing well before determining final pH. Loosen jar cap 1/4 turn and incubate 24 ± 2 h at 35°C . Continue as in D, 1-11, below.
- 9. **Frosting and topping mixes.** Aseptically weigh 25 g sample into sterile, wide-mouth, screw-cap jar (500 ml) or other appropriate container. Add 225 ml [nutrient broth](#) and mix well. Cap jar securely and let stand 60 ± 5 min at room temperature. Mix well by swirling and determine pH with test paper. Adjust pH, if necessary, to 6.8 ± 0.2 . Loosen jar cap about 1/4 turn and incubate 24 ± 2 h at 35°C . Continue as in D, 1-11, below.
- 10. **Spices**
 - a. **Black pepper, white pepper, celery seed or flakes, chili powder, cumin, paprika, parsley flakes, rosemary, sesame seed, thyme, and vegetable flakes.**

Aseptically weigh 25 g sample into sterile, wide-mouth, screw-cap jar (500 ml) or other appropriate container. Add 225 ml sterile [trypticase soy broth \(TSB\)](#) and mix well. Cap jar securely and let stand 60 ± 5 min at room temperature. Mix well by swirling and determine pH with test paper. Adjust pH, if necessary, to 6.8 ± 0.2 . Loosen jar cap about 1/4 turn and incubate 24 ± 2 h at 35°C . Continue as in D, 1-11, below.

- b. **Onion flakes, onion powder, garlic flakes.**

Aseptically weigh 25 g sample into sterile, wide-mouth, screw-cap jar (500 ml) or other appropriate container. Preenrich sample in [TSB](#) with added K_2SO_3 (5 g K_2SO_3 per 1000 ml TSB, resulting in final 0.5% K_2SO_3 concentration). Add K_2SO_3 to broth before autoclaving 225 ml volumes in 500 ml Erlenmeyer flasks at 121°C for 15 min. After autoclaving, aseptically determine and, if necessary, adjust final volume to 225 ml. Add 225 ml sterile TSB with added K_2SO_3 to sample and mix well. Continue as in C-10a.

- c. **Allspice, cinnamon, cloves, and oregano.**

At this time there are no known methods for neutralizing the toxicity of these 4 spices. Dilute them beyond their toxic levels to examine them. Examine allspice, cinnamon, and oregano at 1:100 sample/broth ratio, and cloves at 1:1000 sample/broth ratio. Examine leafy condiments at sample/broth ratio greater than 1:10 because of physical difficulties encountered by absorption of broth by dehydrated product. Examine these spices as described in C-10a, above, maintaining recommended sample/broth ratios.

11. **Candy and candy coating (including chocolate).** Aseptically weigh 25 g sample into sterile blending container. Add 225 ml sterile, [reconstituted nonfat dry milk](#) and blend 2 min. Aseptically transfer homogenized mixture to sterile, wide-mouth, screw-cap jar (500 ml) or other appropriate container and let stand 60 ± 5 min at room temperature with jar securely capped. Mix well by swirling and determine pH with test paper. Adjust pH, if necessary, to 6.8 ± 0.2 . Add 0.45 ml 1% aqueous brilliant green dye solution and mix well. Loosen jar caps 1/4 turn and incubate 24 ± 2 h at 35°C . Continue as in D, 1-11, below.
12. **Coconut.** Aseptically weigh 25 g sample into sterile, wide-mouth, screw-cap jar (500 ml) or other appropriate container. Add 225 ml sterile [lactose broth](#), shake well, and let stand 60 ± 5 min at room temperature with jar securely capped. Mix well by swirling and determine pH with test paper. Adjust pH, if necessary, to 6.8 ± 0.2 . Add up to 2.25 ml steamed (15 min) [Tergitol Anionic 7](#) and mix well. Alternatively, use steamed (15 min) [Triton X-100](#). Limit use of these surfactants to minimum quantity needed to initiate foaming. For Triton X-100 this quantity may be as little as 2 or 3 drops. Loosen jar cap about 1/4 turn and incubate 24 ± 2 h at 35°C . Continue as in D, 1-11, below.
13. **Food dyes and food coloring substances.** For dyes with pH 6.0 or above (10% aqueous suspension), use method described for dried whole eggs (C-1, above). For laked dyes or dyes with pH below 6.0, aseptically weigh 25 g sample into sterile, wide-mouth, screw-cap jar (500 ml) or other appropriate container. Add 225 ml [tetrathionate broth](#) without brilliant green dye. Mix well and let stand 60 ± 5 min at room temperature with jar securely capped. Using pH meter, adjust pH to 6.8 ± 0.2 . Add 2.25 ml [0.1% brilliant green dye solution](#) and mix thoroughly by swirling. Loosen jar cap about 1/4 turn and incubate 24 ± 2 h at 35°C . Continue as in D, 3-11, below.
14. **Gelatin.** Aseptically weigh 25 g sample into sterile, wide-mouth, screw-cap jar (500 ml) or other appropriate container. Add 225 ml sterile [lactose broth](#) and 5 ml 5% aqueous [papain solution](#) and mix well. Cap jar securely and incubate at 35°C for 60 ± 5 min. Mix well by swirling and determine pH with test paper. Adjust pH, if necessary, to 6.8 ± 0.2 . Loosen jar cap about 1/4 turn and incubate 24 ± 2 h at 35°C . Continue as in D, 1-11, below.
15. **Meats, meat substitutes, meat by-products, animal substances, glandular products, and meals (fish, meat, bone).** Aseptically weigh 25 g sample into sterile blending container. Add 225 ml sterile [lactose broth](#) and blend 2 min. Aseptically transfer homogenized mixture to sterile wide-mouth, screw-cap jar (500 ml) or other appropriate container and let stand 60 ± 5 min at room temperature with jar securely capped. If mixture is powder or is ground or comminuted, blending may be omitted. For samples that do not require blending, add lactose broth and mix thoroughly; let stand for 60 ± 5 min at room temperature with jar securely capped.

Mix well by swirling and determine pH with test paper. Adjust pH, if necessary, to 6.8 ± 0.2 . Add up to 2.25 ml steamed (15 min) Tergitol Anionic 7 and mix well. Alternatively, use steamed (15 min) Triton X-100. Limit use of these surfactants to minimum quantity needed to initiate foaming. Actual quantity will depend on composition of test material. Surfactants will not be needed in analysis of powdered glandular products. Loosen jar caps 1/4 turn and incubate sample mixtures 24 ± 2 h at 35°C . Continue as in D, 1-11, below.

16. **Frog legs.** (This method is used for all domestic and imported frog legs.) Place 15 pairs of frog legs into sterile plastic bag and cover with sterile lactose broth at a 1:9 sample-to-

broth (g/ml) ratio (see A, 23-24, above). If single legs are estimated to average 25 g or more, examine only one leg of each of 15 pairs. Place bag in large plastic beaker or other suitable container. Mix well and let stand 60 ± 5 min at room temperature. Mix well by swirling and determine pH with test paper. Adjust pH, if necessary, to 6.8 ± 0.2 . Place plastic bag containing the frog legs and lactose broth into plastic beaker or other suitable container. Incubate 24 ± 2 h at 35°C . Continue examination as in D, 1-11, below.

17. **Rabbit carcasses.** (This method is used for all domestic and imported rabbit carcasses.) Place rabbit carcass into sterile plastic bag. Place bag in beaker or other suitable container. Add sterile lactose broth at a 1:9 sample-to-broth (g/ml) ratio to cover carcass (see A, 23-24, above). Mix well by swirling and let stand 60 ± 5 min at room temperature. Mix well by swirling and determine pH with test paper. Adjust pH, if necessary, to 6.8 ± 0.2 . Incubate 24 ± 2 h at 35°C . Continue examination as in D, 1-11, below.
18. **Guar gum.** Aseptically weigh 25 g sample into sterile beaker (250 ml) or other appropriate container. Prepare a 1.0% cellulase solution (add 1 g cellulase to 99 ml sterile distilled water). Dispense into 150 ml bottles. (Cellulase solution may be stored at $2-5^\circ\text{C}$ for up to 2 weeks). Add 225 ml sterile [lactose broth](#) and 2.25 ml sterile 1% cellulase solution to sterile, wide-mouth, screw-cap jar (500 ml) or other appropriate container. While vigorously stirring the cellulase/lactose broth with magnetic stirrer, pour 25 g analytical unit quickly through sterile glass funnel into the cellulase/lactose broth. Cap jar securely and let stand 60 ± 5 min at room temperature. Incubate loosely capped container without pH adjustment, for 24 ± 2 h at 35°C . Continue as in D, 1-11, below.
19. **Orange juice (pasteurized and unpasteurized), apple cider (pasteurized and unpasteurized), and apple juice (pasteurized)** Aseptically add 25 ml sample to 225 ml [Universal preenrichment broth](#) in a sterile, wide mouth, screw-capped jar (500 ml) or other appropriate container. Swirl the flask contents thoroughly. Cap jar securely and let stand 60 ± 5 min at room temperature. Do not adjust pH. Incubate loosely capped container for 24 ± 2 h at 35°C . Continue as in D, 1-11, below (treat as a low microbial load food).
20. **Pig ears and other types of dog chew pieces.** Place 1 piece (or 2-3 pieces if smaller sizes) from each sample unit into sterile plastic bag. Place bag into large beaker or other suitable container. Add sterile lactose broth at a 1:9 sample-to-broth (g/ml) ratio to cover pieces (see A, 23-24, above). Mix well by swirling and let stand 60 ± 5 min at room temperature. Mix well by swirling and determine pH with test paper. Adjust pH, if necessary, to 6.8 ± 0.2 . Add either steamed (15 min) Tergitol Anionic 7 or steamed (15 min) Triton X-100 up to a 1% concentration. For example, if 225 ml lactose broth is added, the maximum volume of added surfactant is 2.25 ml. Limit use of these surfactants to minimum quantity to initiate foaming. Incubate 24 ± 2 h at 35°C . Continue examination as in D, 1-11, below.
21. **Cantaloupes.** Preferably, do not thaw frozen samples before analysis. If frozen sample must be tempered to obtain analytical portion, thaw below 45°C for <15 min with continuous agitation in thermostatically controlled water bath or thaw within 18 h at $2-5^\circ\text{C}$.

For comminuted or cut fruit, aseptically weigh 25 g sample into sterile blending container. Add 225 ml sterile [Universal preenrichment broth](#) (UP) and blend 2 min. Aseptically transfer homogenized mixture to sterile, wide-mouth, screw-cap jar (500 ml) or other appropriate container and let stand 60 ± 5 min at room temperature with jar securely capped. Do not adjust pH. Mix well and loosen jar cap about 1/4 turn. Incubate 24 ± 2 h at 35°C . Continue as in D, 1-11, below.

For whole cantaloupes, do not rinse even if there is visible dirt. Examine the cantaloupes "as is".

Place the cantaloupe into a sterile plastic bag. Add enough [UP](#) broth to allow the cantaloupe to float. The volume of [UP](#) broth may be 1.5 times the weight of the cantaloupes. For example, cantaloupes weighing 1500 g will probably need a volume of approximately 2250 ml [UP](#) broth to float. Add more broth, if necessary. Place the plastic bag, with cantaloupes and [UP](#) broth, into a 5 liter beaker, or other appropriate container,

for support during incubation. Allow the open-end flap of the plastic bag to "fold over" so as to form a secure, but not air-tight, closure during incubation.

Let stand for 60 ± 5 min at room temperature. Do not adjust pH. Incubate slightly opened bag, containing cantaloupe, for 24 ± 2 h at 35°C . Continue as in D, 1-11, below.

22. **Mangoes.** Preferably, do not thaw frozen samples before analysis. If frozen sample must be tempered to obtain analytical portion, thaw below 45°C for <15 min with continuous agitation in thermostatically controlled water bath or thaw within 18 h at $2-5^{\circ}\text{C}$.

For comminuted or cut fruit, aseptically weigh 25 g sample into sterile blending container. Add 225 ml sterile [buffered peptone water \(BPW\)](#) and blend 2 min. Aseptically transfer homogenized mixture to sterile, wide-mouth, screw-cap jar (500 ml) or other appropriate container and let stand 60 ± 5 min at room temperature with jar securely capped. Mix well by swirling and determine pH with test paper. Adjust pH, if necessary, to 6.8 ± 0.2 . Mix well and loosen jar cap about 1/4 turn. Incubate 24 ± 2 h at 35°C . Continue as in D, 1-11, below.

For whole mangoes, do not rinse even if there is visible dirt. Examine the mangoes "as is".

Place the mango into a sterile plastic bag. Add enough [BPW](#) to allow the mango to float. The volume of [BPW](#) may be 1.0 times the weight of the mangoes. For example, mangoes weighing 500 g will probably need a volume of approximately 500 ml [BPW](#) broth to float. Add more broth, if necessary. Place the plastic bag, with mangoes and [BPW](#) broth, into a 5 liter beaker, or other appropriate container, for support during incubation.

Let stand for 60 ± 5 min at room temperature. Adjust pH to 6.8 ± 0.2 , if necessary. Incubate slightly opened bag for 24 ± 2 h at 35°C . Continue as in D, 1-11, below.

23. **Tomatoes.** For comminuted or cut fruit, aseptically weigh 25 g sample into sterile blending container. Add 225 ml sterile buffered peptone water and blend 2 min. Aseptically transfer homogenized mixture to sterile, wide-mouth, screw-cap jar (500 ml) or other appropriate container and let stand 60 ± 5 min at room temperature with jar securely capped. Mix well by swirling and determine pH with test paper. Adjust pH, if necessary, to 6.8 ± 0.2 . Mix well and loosen jar cap about 1/4 turn. Incubate 24 ± 2 h at 35°C . Continue as in D, 1-11, below.

For whole tomatoes, do not rinse even if there is visible dirt. Examine the tomatoes "as is".

Place the tomato into a sterile plastic bag or other suitable container (sterile foil covered beaker can be used). Add enough [UP](#) broth to allow the tomato to float. The volume of [UP](#) broth may be 1.0 times the weight of the tomato. For example, tomatoes weighing 300 g will probably need a volume of approximately 300 ml [UP](#) broth to float. Add more, if necessary. Place the plastic bag (if used), with tomato and [UP](#) broth, into a sterile beaker (beaker size is dependent on the size of the tomato), or other appropriate container, for support during incubation. Allow the open-end flap of the plastic bag to "fold over" so as to form a secure, but not air-tight, closure during incubation.

Let stand for 60 ± 5 min at room temperature. Do not adjust pH. Incubate slightly opened bag for 24 ± 2 h at 35°C . Continue as in D, 1-11, below.

24. **Environmental testing.** Sample environmental surfaces with sterile swabs or sponges. Place the swab/sponge in a sterile Whirl-pak bag, or equivalent, that contains enough [Dey-Engley \(DE\) broth](#) to cover the swab/sponge.

Transport swabs/sponges in an insulated transport container with frozen gel packs to keep the samples cold, but not frozen. If samples cannot be processed immediately, refrigerate at $4 \pm 2^{\circ}\text{C}$. Start sample analysis within 48 ± 2 h of collection.

Add swab/sponge to 225 ml lactose broth in a sterile, wide mouth, screw-capped jar (500 ml) or other appropriate container. Swirl the flask contents thoroughly. Cap jar securely and let stand 60 ± 5 min at room temperature. Mix well by swirling and determine pH with test paper. Adjust pH, if necessary, to 6.8 ± 0.2 . Incubate 24 ± 2 h at 35°C . Continue examination as in D, 1-11, below.

25. **Alfalfa seeds and mung beans.** Aseptically weigh 25g alfalfa seeds or mung beans into a sterile 500 mL Erlenmeyer flask. Aseptically add 225 mL lactose broth to the test portion and swirl the Erlenmeyer flask. Cover the mouth of the Erlenmeyer flask with sterile aluminum foil and allow contents to stand at room temperature for 60 ± 5 min. Adjust the pH of the culture to 6.8 ± 0.2 , if necessary. Incubate for 24 ± 2 h at $35 \pm 2^{\circ}\text{C}$. Continue as in D, 1-11, below (treat as high microbial load food).
26. **Mamey pulp.** If frozen, sample must be tempered to obtain analytical portion. Thaw below 45°C for <15 min with continuous agitation in thermostatically controlled water bath or thaw within 18 h at $2-5^{\circ}\text{C}$.

For mamey pulp, suspected to be contaminated with *S. Typhi*, aseptically weigh 25 g sample into sterile, wide-mouth, screw-cap jar (500 ml) or other appropriate container. Add 225 ml sterile [Universal Preenrichment broth without ferric ammonium citrate](#), mix by swirling, and let stand 60 ± 5 min at room temperature with jar securely capped. Do not adjust pH. Mix well and loosen jar cap about 1/4 turn. Incubate 24 ± 2 h at 35°C . Continue as in D, 1-11, below. Treat as a low microbial load food.

For mamey pulp, NOT suspected to be contaminated with *S. Typhi*, aseptically weigh 25 g sample into sterile, wide-mouth, screw-cap jar (500 ml) or other appropriate container. Add 225 ml sterile Universal Preenrichment broth, mix by swirling, and let stand 60 ± 5 min at room temperature with jar securely capped. Do not adjust pH. Mix well and loosen jar cap about 1/4 turn. Incubate 24 ± 2 h at 35°C . Continue as in D, 1-11, below.

[Return to Chapter Contents](#)

D. [Isolation of *Salmonella*](#)

1. Tighten lid and gently shake incubated sample.

Guar gum and foods suspected to be contaminated with *S. Typhi*. Transfer 1 ml mixture to 10 ml [selenite cystine \(SC\) broth](#) and another 1 ml mixture to 10 ml [TT broth](#)⁹⁸. Vortex.

All other foods. Transfer 0.1 ml mixture to 10 ml [Rappaport-Vassiliadis \(RV\) medium](#) and another 1 ml mixture to 10 ml [tetrathionate \(TT\) broth](#). Vortex.

2. Incubate selective enrichment media as follows:

Foods with a high microbial load. Incubate RV medium 24 ± 2 h at $42 \pm 0.2^{\circ}\text{C}$ (circulating, thermostatically-controlled, water bath). Incubate TT broth 24 ± 2 h at $43 \pm 0.2^{\circ}\text{C}$ (circulating, thermostatically-controlled, water bath).

Foods with a low microbial load (except guar gum and foods suspected to be contaminated with *S. Typhi*). Incubate RV medium 24 ± 2 h at $42 \pm 0.2^\circ\text{C}$ (circulating, thermostatically controlled, water bath). Incubate TT broth 24 ± 2 h at $35 \pm 2.0^\circ\text{C}$.

Guar gum and foods suspected to be contaminated with *S. Typhi*. Incubate SC and TT broths 24 ± 2 h at 35°C .

3. Mix (vortex, if tube) and streak 3 mm loopful (10 μl) incubated TT broth on [bismuth sulfite \(BS\) agar](#), [xylose lysine desoxycholate \(XLD\) agar](#), and [Hektoen enteric \(HE\) agar](#). **Prepare BS plates the day before streaking and store in dark at room temperature until streaked.**
4. Repeat with 3 mm loopful (10 μl) of RV medium (for samples of high and low microbial load foods) and of SC broth (for guar gum).
5. Refer to 994.04 in *Official Methods of Analysis* (1) for option of refrigerating incubated sample preenrichments and incubated sample selective enrichments (SC and TT broths only) of low moisture foods. This option allows sample analyses to be initiated as late as Thursday while still avoiding weekend work.
6. Incubate plates 24 ± 2 h at 35°C .
7. Examine plates for presence of colonies that may be *Salmonella*.

TYPICAL *Salmonella* COLONY MORPHOLOGY

Pick 2 or more colonies of *Salmonella* from each selective agar after 24 ± 2 h incubation. Typical *Salmonella* colonies are as follows:

- a. **Hektoen enteric (HE) agar.** Blue-green to blue colonies with or without black centers. Many cultures of *Salmonella* may produce colonies with large, glossy black centers or may appear as almost completely black colonies.
- b. **Xylose lysine desoxycholate (XLD) agar.** Pink colonies with or without black centers. Many cultures of *Salmonella* may produce colonies with large, glossy black centers or may appear as almost completely black colonies.
- c. **Bismuth sulfite (BS) agar.** Brown, gray, or black colonies; sometimes they have a metallic sheen. Surrounding medium is usually brown at first, but may turn black in time with increased incubation, producing the so-called halo effect.

If typical colonies are present on the BS agar after 24 ± 2 h incubation, then pick 2 or more colonies. Irrespective of whether or not BS agar plates are picked at 24 ± 2 h, reincubate BS agar plates an additional 24 ± 2 h. After 48 ± 2 h incubation, pick 2 or more typical colonies, if present, from the BS agar plates, only if colonies picked from the BS agar plates incubated for 24 ± 2 h give atypical reactions in triple sugar iron agar (TSI) and lysine iron agar (LIA) that result in culture being discarded as not being *Salmonella*. See sections D.9 and D.10, below, for details in interpreting TSI and LIA reactions.

ATYPICAL *Salmonella* COLONY MORPHOLOGY

In the absence of typical or suspicious *Salmonella* colonies, search for atypical *Salmonella* colonies as follows:

- d. **HE and XLD agars.** Atypically a few *Salmonella* cultures produce yellow colonies with or without black centers on HE and XLD agars. In the absence of typical *Salmonella* colonies on HE or XLD agars after 24 ± 2 h incubation, then pick 2 or more atypical *Salmonella* colonies.
- e. **BS agar.** Atypically some strains produce green colonies with little or no darkening of the surrounding medium. If typical or suspicious colonies are not

present on BS agar after 24 ± 2 h, then do not pick any colonies but reincubate an additional 24 ± 2 h. If typical or suspicious colonies are not present after 48 ± 2 h incubation, then pick 2 or more atypical colonies.

SUGGESTED CONTROL CULTURES

In addition to the positive control cultures (typical *Salmonella*), 3 additional *Salmonella* cultures are recommended to assist in the selection of atypical *Salmonella* colony morphology on selective agars. These cultures are a lactose-positive, H₂S-positive *S. diarizonae* (ATCC 12325) and a lactose-negative, H₂S-negative *S. abortus equi* (ATCC 9842); **OR** a lactose-positive, H₂S-negative *S. diarizonae* (ATCC 29934). These cultures may be obtained from the [American Type Culture Collection](#), 10801 University Boulevard, Manassas, VA 20110-2209.

8. Lightly touch the very center of the colony to be picked with sterile inoculating needle and inoculate TSI slant by streaking slant and stabbing butt. Without flaming, inoculate LIA slant by stabbing butt twice and then streaking slant. Since lysine decarboxylation reaction is strictly anaerobic, the LIA slants must have deep butt (4 cm). Store picked selective agar plates at 5-8°C.
9. Incubate TSI and LIA slants at 35°C for 24 ± 2 h. Cap tubes loosely to maintain aerobic conditions while incubating slants to prevent excessive H₂S production. *Salmonella* in culture typically produces alkaline (red) slant and acid (yellow) butt, with or without production of H₂S (blackening of agar) in TSI. In LIA, *Salmonella* typically produces alkaline (purple) reaction in butt of tube. Consider only distinct yellow in butt of tube as acidic (negative) reaction. Do not eliminate cultures that produce discoloration in butt of tube solely on this basis. Most *Salmonella* cultures produce H₂S in LIA. Some non-*Salmonella* cultures produce a brick-red reaction in LIA slants.
10. All cultures that give an alkaline butt in LIA, regardless of TSI reaction, should be retained as potential *Salmonella* isolates and submitted for biochemical and serological tests. Cultures that give an acid butt in LIA and an alkaline slant and acid butt in TSI should also be considered potential *Salmonella* isolates and should be submitted for biochemical and serological tests. Cultures that give an acid butt in LIA and an acid slant and acid butt in TSI may be discarded as not being *Salmonella*. Test retained, presumed-positive TSI cultures as directed in D-11, below, to determine if they are *Salmonella* species, including *S. arizonae*. If TSI cultures fail to give typical reactions for *Salmonella* (alkaline slant and acid butt) pick additional suspicious colonies from selective medium plate not giving presumed-positive culture and inoculate TSI and LIA slants as described in D-8, above.
11. Apply biochemical and serological identification tests to:
 - a. Three presumptive TSI cultures recovered from set of plates streaked from RV medium (or SC broth for guar gum), if present, and 3 presumptive TSI agar cultures recovered from plates streaked from TT broth, if present.
 - b. If 3 presumptive-positive TSI cultures are not isolated from one set of agar plates, test other presumptive-positive TSI agar cultures, if isolated, by biochemical and serological tests. Examine a minimum of 6 TSI cultures for each 25 g analytical unit or each 375 g composite.

Return to Chapter Contents

E. Identification of *Salmonella*

1. **Mixed cultures.** Streak TSI agar cultures that appear to be mixed on [MacConkey agar](#), [HE agar](#), or [XLD agar](#). Incubate plates 24 ± 2 h at 35°C. Examine plates for presence of colonies suspected to be *Salmonella*.
 - a. **MacConkey agar.** Typical colonies appear transparent and colorless, sometimes with dark center. Colonies of *Salmonella* will clear areas of precipitated bile caused by other organisms sometimes present.

- b. **Hektoen enteric (HE) agar.** See D-7a, above.
 - c. **Xylose lysine desoxycholate (XLD) agar.** See D-7b, above. Transfer at least 2 colonies suspected to be *Salmonella* to TSI and LIA slants as described in D-7, above, and continue as in D-9, above.
2. **Pure cultures**
- a. **Urease test (conventional).** With sterile needle, inoculate growth from each presumed-positive TSI slant culture into tubes of [urea broth](#). Since occasional, uninoculated tubes of urea broth turn purple-red (positive test) on standing, include uninoculated tube of this broth as control. Incubate 24 ± 2 h at 35°C .
 - b. **Optional urease test (rapid).** Transfer two 3-mm loopfuls of growth from each presumed-positive TSI slant culture into tubes of [rapid urea broth](#). Incubate 2 h in $37 \pm 0.5^\circ\text{C}$ water bath. Discard all cultures giving positive test. Retain for further study all cultures that give negative test (no change in color of medium).
3. **Serological polyvalent flagellar (H) test**
- a. Perform the polyvalent flagellar (H) test at this point, or later, as described in E-5, below. Inoculate growth from each urease-negative TSI agar slant into either 1) [BHI broth](#) and incubate 4-6 h at 35°C until visible growth occurs (to test on same day); or 2) [trypticase soy-tryptose broth](#) and incubate 24 ± 2 h at 35°C (to test on following day). Add 2.5 ml formalinized physiological saline solution to 5 ml of either broth culture.
 - b. Select 2 formalinized broth cultures and test with *Salmonella* polyvalent flagellar (H) antisera. Place 0.5 ml of appropriately diluted *Salmonella* polyvalent flagellar (H) antiserum in 10 x 75 mm or 13 x 100 mm serological test tube. Add 0.5 ml antigen to be tested. Prepare saline control by mixing 0.5 ml formalinized physiological saline solution with 0.5 ml formalinized antigen. Incubate mixtures in $48\text{-}50^\circ\text{C}$ water bath. Observe at 15 min intervals and read final results in 1 h.

Positive--agglutination in test mixture and no agglutination in control.

Negative--no agglutination in test mixture and no agglutination in control.

Nonspecific--agglutination in both test mixture and control. Test the cultures giving such results with Spicer-Edwards antisera.

4. **Spicer-Edwards serological test.** Use this test as an alternative to the polyvalent flagellar (H) test. It may also be used with cultures giving nonspecific agglutination in polyvalent flagellar (H) test. Perform Spicer-Edwards flagellar (H) antisera test as described in E, 3b, above. Perform additional biochemical tests (E, 5a-c, below) on cultures giving positive flagellar test results. If both formalinized broth cultures are negative, perform serological tests on 4 additional broth cultures (E, 3a, above). If possible, obtain 2 positive cultures for additional biochemical testing E, 5a-c, below). If all urease-negative TSI cultures from sample give negative serological flagellar (H) test results, perform additional biochemical tests E, 5a-c, below).
5. **Testing of urease-negative cultures**
- a. [Lysine decarboxylase broth](#). If LIA test was satisfactory, it need not be repeated. Use lysine decarboxylase broth for final determination of lysine decarboxylase if culture gives doubtful LIA reaction. Inoculate broth with small amount of growth from TSI slant suspicious for *Salmonella*. Replace cap tightly and incubate 48 ± 2 h at 35°C but examine at 24 h intervals. *Salmonella* species cause alkaline reaction indicated by purple color throughout medium. Negative test is indicated by yellow color throughout medium. If medium appears discolored (neither purple nor yellow) add a few drops of 0.2% bromocresol purple dye and re-read tube reactions.
 - b. [Phenol red dulcitol broth](#) or [purple broth base with 0.5% dulcitol](#). Inoculate broth with small amount of growth from TSI culture. Replace cap loosely and incubate 48 ± 2 h at 35°C , but examine after 24 h. Most *Salmonella* species give positive test, indicated by gas formation in inner fermentation vial and acid pH

(yellow) of medium. Production of acid should be interpreted as a positive reaction. Negative test is indicated by no gas formation in inner fermentation vial and red (with phenol red as indicator) or purple (with bromocresol purple as indicator) color throughout medium.

- c. **Tryptone (or tryptophane) broth.** Inoculate broth with small growth from TSI agar culture. Incubate 24 ± 2 h at 35°C and proceed as follows:
 1. **1) Potassium cyanide (KCN) broth.** Transfer 3 mm loopful of 24 h tryptophane broth culture to KCN broth. Heat rim of tube so that good seal is formed when tube is stoppered with wax-coated cork. Incubate 48 ± 2 h at 35°C but examine after 24 h. Interpret growth (indicated by turbidity) as positive. Most *Salmonella* species do not grow in this medium, as indicated by lack of turbidity.
 2. **2) Malonate broth.** Transfer 3 mm loopful of 24 h tryptone broth culture to malonate broth. Since occasional uninoculated tubes of malonate broth turn blue (positive test) on standing, include uninoculated tube of this broth as control. Incubate 48 ± 2 h at 35°C , but examine after 24 h. Most *Salmonella* species cultures give negative test (green or unchanged color) in this broth.
 3. **3) Indole test.** Transfer 5 ml of 24 h tryptophane broth culture to empty test tube. Add 0.2-0.3 ml Kovacs' reagent. Most *Salmonella* cultures give negative test (lack of deep red color at surface of broth). Record intermediate shades of orange and pink as \pm .
 4. **4) Serological flagellar (H) tests for *Salmonella*.** If either polyvalent flagellar (H) test (E-3, above) or the Spicer-Edwards flagellar (H) test tube test (E-4, above) has not already been performed, either test may be performed here.
 5. **5) Discard as not *Salmonella*** any culture that shows either positive indole test and negative serological flagellar (H) test, or positive KCN test and negative lysine decarboxylase test.
6. **Serological somatic (O) tests for *Salmonella*.**
(Pre-test all antisera to *Salmonella* with known cultures.)
 - a. **Polyvalent somatic (O) test.** Using wax pencil, mark off 2 sections about 1 x 2 cm each on inside of glass or plastic petri dish (15 x 100 mm). Commercially available sectioned slides may be used. Emulsify 3 mm loopful of culture from 24-48 h TSI slant or, preferably, tryptose blood agar base (without blood) with 2 ml 0.85% saline. Add 1 drop of culture suspension to upper portion of each rectangular crayon-marked section. Add 1 drop of saline solution to lower part of one section only. Add 1 drop of *Salmonella* polyvalent somatic (O) antiserum to other section only. With clean sterile transfer loop or needle, mix culture suspension with saline solution for one section and repeat for other section containing antiserum. Tilt mixtures in back-and-forth motion for 1 min and observe against dark background in good illumination. Consider any degree of agglutination a positive reaction. Classify polyvalent somatic (O) test results as follows:

Positive--agglutination in test mixture; no agglutination in saline control.

Negative--no agglutination in test mixture; no agglutination in saline control.

Nonspecific--agglutination in test and in control mixtures. Perform further biochemical and serological tests as described in *Edwards and Ewing's Identification of Enterobacteriaceae* (2).
 - b. **Somatic (O) group tests.** Test as in E-6a, above, using individual group somatic (O) antisera including Vi, if available, in place of *Salmonella* polyvalent somatic (O) antiserum. For special treatment of cultures giving positive Vi agglutination reaction, refer to sec. 967.28B in *Official Methods of Analysis* (1). Record

cultures that give positive agglutination with individual somatic (O) antiserum as positive for that group. Record cultures that do not react with individual somatic (O) antiserum as negative for that group.

7. **Additional biochemical tests.** Classify as *Salmonella* those cultures which exhibit typical *Salmonella* reactions for tests 1-11, shown in Table 1. If one TSI culture from 25 g analytical unit is classified as *Salmonella*, further testing of other TSI cultures from the same 25 g analytical unit is unnecessary. Cultures that contain demonstrable *Salmonella* antigens as shown by positive *Salmonella* flagellar (H) test but do not have biochemical characteristics of *Salmonella* should be purified (E-1, above) and retested, beginning with E-2, above.

Perform the following additional tests on cultures that do not give typical *Salmonella* reactions for tests 1-11 in Table 1 and that consequently do not classify as *Salmonella*.

a. **Phenol red lactose broth or purple lactose broth.**

- **1)** Inoculate broth with small amount of growth from unclassified 24-48 h TSI slant. Incubate 48 ± 2 h at 35°C , but examine after 24 h.

Positive--acid production (yellow) and gas production in inner fermentation vial. Consider production of acid only as positive reaction. Most cultures of *Salmonella* give negative test result, indicated by no gas formation in inner fermentation vial and red (with phenol red as indicator) or purple (with bromocresol purple as indicator) throughout medium.

- **2)** Discard as not *Salmonella*, cultures that give positive lactose tests, except cultures that give acid slants in TSI and positive reactions in LIA, or cultures that give positive malonate broth reactions. Perform further tests on these cultures to determine if they are *S. arizonae*.

b. **Phenol red sucrose broth or purple sucrose broth.** Follow procedure described in E,7a-1, above. Discard as not *Salmonella*, cultures that give positive sucrose tests, except those that give acid slants in TSI and positive reactions in LIA.

c. **MR-VP broth.** Inoculate medium with small amount of growth from each unclassified TSI slant suspected to contain *Salmonella*. Incubate 48 ± 2 h at 35°C .

- **1)** Perform Voges-Proskauer (VP) test at room temperature as follows: Transfer 1 ml 48 h culture to test tube and incubate remainder of MR-VP broth an additional 48 h at 35°C . Add 0.6 ml α -naphthol and shake well. Add 0.2 ml 40% KOH solution and shake. To intensify and speed reaction, add a few crystals of creatine. Read results after 4 h; development of pink-to-ruby red color throughout medium is positive test. Most cultures of *Salmonella* are VP-negative, indicated by absence of development of pink-to-red color throughout broth.
- **2)** Perform methyl red test as follows: To 5 ml of 96 h MR-VP broth, add 5-6 drops of methyl red indicator. Read results immediately. Most *Salmonella* cultures give positive test, indicated by diffuse red color in medium. A distinct yellow color is negative test. Discard, as not *Salmonella*, cultures that give positive KCN and VP tests and negative methyl red test.

d. **Simmons citrate agar.** Inoculate this agar, using needle containing growth from unclassified TSI agar slant. Inoculate by streaking slant and stabbing butt. Incubate 96 ± 2 h at 35°C . Read results as follows:

Positive--presence of growth, usually accompanied by color change from green to blue. Most cultures of *Salmonella* are citrate-positive.

Negative--no growth or very little growth and no color change.

8. **Classification of cultures.** Classify, as *Salmonella*, cultures that have reaction patterns of Table I. Discard, as not *Salmonella*, cultures that give results listed in any subdivision of Table 2. Perform additional tests described in *Edwards and Ewing's Identification of Enterobacteriaceae (2)* to classify any culture that is not clearly identified as *Salmonella* by classification scheme in Table I or not eliminated as not being *Salmonella* by test reactions in Table 2. If neither of 2 TSI cultures carried through biochemical tests confirms the isolate as *Salmonella*, perform biochemical tests, beginning with E-5, on remaining urease-negative TSI cultures from same 25 g analytical unit.

Table 1. Biochemical and serological reactions of <i>Salmonella</i>			
Test or substrate	Result		<i>Salmonella</i> species reaction ^(a)
	Positive	Negative	
1. Glucose (TSI)	yellow butt	red butt	+
2. Lysine decarboxylase (LIA)	purple butt	yellow butt	+
3. H ₂ S (TSI and LIA)	Blackening	no blackening	+
4. Urease	purple-red color	no color change	-
5. Lysine decarboxylase broth	purple color	yellow color	+
6. Phenol red dulcitol broth	yellow color and/or gas	no gas; no color change	+ ^(b)
7. KCN broth	Growth	no growth	-
8. Malonate broth	blue color	no color change	- ^(c)
9. Indole test	violet color at surface	yellow color at surface	-
10. Polyvalent flagellar test	Agglutination	no agglutination	+
11. Polyvalent somatic test	Agglutination	no agglutination	+
12. Phenol red lactose broth	yellow color and/or gas	no gas; no color change	- ^(c)
13. Phenol red sucrose broth	yellow color and/or gas	no gas; no color change	-
14. Voges-Proskauer test	pink-to-red color	no color change	-
15. Methyl red test	diffuse red color	diffuse yellow color	+
16. Simmons citrate	growth; blue color	no growth; no color change	v

^a +, 90% or more positive in 1 or 2 days; -, 90% or more negative in 1 or 2 days; v, variable.

^b Majority of *S. arizonae* cultures are negative.

^c Majority of *S. arizonae* cultures are positive.

Table 2. Criteria for discarding non- <i>Salmonella</i> cultures	
Test or substrate	Results
1. Urease	positive (purple-red color)
2. Indole test and Polyvalent flagellar (H) test;	positive (violet color at surface) negative (no agglutination)
or Indole test and Spicer-Edwards flagellar test	positive (violet color at surface) negative (no agglutination)
3. Lysine decarboxylase and KCN broth	negative (yellow color) positive (growth)
4. Phenol red lactose broth	positive (yellow color and/or gas) ^{(a), (b)}
5. Phenol red sucrose broth	positive (yellow color and/or gas) ^(b)
6. KCN broth, Voges-Proskauer test, and Methyl red test	positive (growth) positive (pink-to-red color) negative (diffuse yellow color)
<p>^a Test malonate broth positive cultures further to determine if they are <i>S. arizonae</i>.</p> <p>^b Do not discard positive broth cultures if corresponding LIA cultures give typical <i>Salmonella</i> reactions; test further to determine if they are <i>Salmonella</i> species.</p>	

9. **Presumptive generic identification of *Salmonella***. As alternative to conventional biochemical tube system, use any of 5 commercial biochemical systems (API 20E, Enterotube II, *Enterobacteriaceae* II, MICRO-ID, or Vitek GNI) for presumptive generic identification of foodborne *Salmonella*. Choose a commercial system based on a demonstration in analyst's own laboratory of adequate correlation between commercial system and biochemical tube system delineated in this identification section. Commercial biochemical kits should not be used as a substitute for serological tests (I). Assemble supplies and prepare reagents required for the kit. Inoculate each unit according to Method 978.24 (API 20E, Enterotube II, and *Enterobacteriaceae* II), sec. 989.12 (MICRO-ID), and Method 991.13 (Vitek GNI) in *Official Methods of Analysis* (1), incubating for time and temperature specified. Add reagents, observe, and record results. For presumptive identification, classify cultures, according to ref. 1, above, as *Salmonella* or not *Salmonella*.

For confirmation of cultures presumptively identified as *Salmonella*, perform the *Salmonella* serological somatic (O) test (E-6, above) and the *Salmonella* serological flagellar (H) test (E-3, above) or the Spicer-Edwards flagellar (H) test (E-4, above), and classify cultures according to the following guidelines:

- a. Report as *Salmonella* those cultures classified as presumptive *Salmonella* with commercial biochemical kits when the culture demonstrates positive *Salmonella* somatic (O) test and positive *Salmonella* (H) test.

- b. Discard cultures presumptively classified as not *Salmonella* with commercial biochemical kits when cultures conform to AOAC criteria (1) for classifying cultures as not *Salmonella*.
 - c. For cultures that do not conform to a or b, classify according to additional tests specified in E, 2-7, above, or additional tests as specified by Ewing (2), or send to reference typing laboratory for definitive serotyping and identification.
10. **Treatment of cultures giving negative flagellar (H) test.** If biochemical reactions of certain flagellar (H)-negative culture strongly suggest that it is *Salmonella*, the negative flagellar agglutination may be the result of nonmotile organisms or insufficient development of flagellar antigen. Proceed as follows: Inoculate motility test medium in petri dish, using small amount of growth from TSI slant. Inoculate by stabbing medium once about 10 mm from edge of plate to depth of 2-3 mm. Do not stab to bottom of plate or inoculate any other portion. Incubate 24 h at 35°C. If organisms have migrated 40 mm or more, retest as follows: Transfer 3 mm loopful of growth that migrated farthest to trypticase soy-tryptose broth. Repeat either polyvalent flagellar (H) (E-3, above) or Spicer-Edwards (E-4, above) serological tests. If cultures are not motile after the first 24 h, incubate an additional 24 h at 35°C; if still not motile, incubate up to 5 days at 25°C. Classify culture as nonmotile if above tests are still negative. If flagellar (H)-negative culture is suspected of being a species of *Salmonella* on the basis of its biochemical reactions, FDA laboratories should submit the culture to

FDA Denver Laboratory
Attention Sample Custodian
Denver Federal Center, Building 20
6th Avenue & Kipling Streets
Denver, CO 80225-0087

(Above address effective October 1, 2004)

for further identification and/or serotyping. Laboratories other than FDA should make arrangements with a reference laboratory for the serotyping of *Salmonella* cultures.

11. **Submission of cultures for serotyping.** Submit 1 isolate of each somatic group recovered from each analytical unit, unless otherwise instructed. Submit cultures on BHI agar slants in screw-cap tubes (13 x 100 mm or 16 x 125 mm) with caps secured tightly. Label each tube with sample number, subsample (analytical unit) number, and code, if applicable. Submit a copy of the Collection Report, FD-464, or Import Sample Report, FD-784 for each sample. Place cultures in culture container with official FDA seal. Place accompanying records (E-11, above) inside shipping carton but not within officially sealed culture container. Submit memo or cover letter for each sample number to expedite reporting of results. Prepare cultures for shipment according to requirements for shipment of etiological agents (3). Label secondary shipping container according to ref. 4. Send container by most rapid mail service available. Maintain duplicate cultures of those submitted for serotyping only on those samples under consideration for legal action.

Microbiology Field laboratories should follow the following guidance in sending *Salmonella* isolates for serotyping:

Isolates from NRL, WEAC, SRL and ARL will be serotyped in ARL:

Arkansas Regional Laboratory
3900 NCTR Road Building 26
Jefferson, AR 72079
Attention: Gwendolyn Anderson
Tel # 870-543-4621
Fax# 870-543-4041

Isolates from SAN, PRL-NW, PRL-SW and DEN will be serotyped in DEN

Denver District Laboratory
6th Avenue & Kipling Street
DFC Building 20
Denver, CO 80225-0087
Attention: Doris Farmer
Tel # 303-236-9604
Fax # 303-236-9675

[Return to Chapter Contents](#)

References

1. AOAC INTERNATIONAL. 2000. Official Methods of Analysis, 17th ed., Methods 967.25-967.28, 978.24, 989.12, 991.13, 994.04, and 995.20. AOAC INTERNATIONAL, Gaithersburg, MD.
2. Ewing, W.H. 1986. Edwards and Ewing's Identification of Enterobacteriaceae, 4th ed. Elsevier, New York.
3. Federal Register. 1971. 36(93):8815 (secs d, e, and f).
4. Federal Register. 1972. 37(191):20556 (sec. 173.388(a)).
5. June, G.A., P.S. Sherrod, T.S. Hammack, R.M. Amaguana, and W.H. Andrews. 1995. Relative effectiveness of selenite cystine broth, tetrathionate broth, and Rappaport-Vassiliadis medium for the recovery of *Salmonella* from raw flesh and other highly contaminated foods: Precollaborative study. J. AOAC Int. 78:375-380.
6. Hammack, T.S., R.M. Amaguana, G.A. June, P.S. Sherrod, and W.H. Andrews. 1999. Relative effectiveness of selenite cystine broth, tetrathionate broth, and Rappaport-Vassiliadis medium for the recovery of *Salmonella* from foods with a low microbial load. J. Food Prot. 62:16-21.
7. June, G.A., P.S. Sherrod, T.S. Hammack, R.M. Amaguana, and W.H. Andrews. 1996. Relative effectiveness of selenite cystine broth, tetrathionate broth, and Rappaport-Vassiliadis medium for the recovery of *Salmonella* from raw flesh, highly contaminated foods, and poultry feed: Collaborative study. J. AOAC Int. 79:1307-1323.
8. Hammack, T.S., R.M. Amaguana, and W.H. Andrews. 2000. Rappaport-Vassiliadis medium for the recovery of *Salmonella* from foods with a low microbial load: Collaborative study. J. AOAC Int. (in press).
9. Sherrod, P.S., R.M. Amaguana, W.H. Andrews, G.A. June, and T.S. Hammack. 1995. Relative effectiveness of selective plating agars for the recovery of *Salmonella* species from selected high-moisture foods. J. AOAC Int. 78:679-690.

[Bad Bug Book:](#)
[Salmonella](#)

Chapter 5 Contents

Hypertext Source: Bacteriological Analytical Manual, 8th Edition, Revision A, 1998. Chapter 5.

Authors: Wallace H. Andrews and Thomas.Hammack@fda.hhs.gov

Revisions: 1999-DEC, 2000-MAR, and 2000-AUG Final revision on 2000-NOV-14 (see the Introduction for a summary of changes).

October 25, 2001 - Extension of the applicability of the orange juice method in section C.19 to apple juice and apple cider.

April 2003 - Frog legs method, Lactic casein, Rennet casein, Sodium caseinate and Rabbit carcass methods revised, top ears and other dog chew toys added. Removed section A.25, Mechanical shaker.

June 2006 Edition - Eggs method revised for shell eggs and liquid whole eggs.

December 2007 - Mamey pulp method added, and Section D revised.

To obtain a copy of a prior version not currently posted, please contact Frederick.Fry@fda.hhs.gov

BAM: Rapid Methods for Detecting Foodborne Pathogens

January 2001

Bacteriological Analytical Manual Appendix 1 Rapid Methods for Detecting Foodborne Pathogens

Authors

Introduction

Authors Note: This section differs from others in this manual in that it lists methods that are not necessarily used by FDA. In addition, the detailed protocols for these methods are not presented, and the user is referred to the instructions that accompany the test kits. One reason for this departure is the incremental rate of change and innovation in rapid testing technology. The best of these new techniques should be evaluated individually by user labs for their particular needs, and also collaboratively for possible adoption as official methods by the AOAC International (1).

The following text and tables list many of the commercially available rapid methods; they are classified by the principles underlying the procedure used. The assay principles and some of the detailed procedures are discussed in other chapters of this manual and/or in the literature cited in the tables. The AOAC status of rapid tests is indicated for those methods that have been validated or evaluated by AOAC (1) and have been adopted as AOAC Official methods. However, these methods continue to be modified or adapted, so that published information may not be the most current. Rapid methods are generally used as screening techniques, with negative results accepted as is, but positive results requiring **confirmation** by the appropriate official method, which, in many instances, is cultural. In many other instances, the rapid method has not been validated; therefore, the listing of a method or kit in this chapter in no way constitutes FDA recommendation or approval.

Rapid Methods

The rapid detection of pathogens and other microbial contaminants in food is critical for ensuring the safety of consumers. Traditional methods to detect foodborne bacteria often rely on time-consuming growth in culture media, followed by isolation, biochemical identification, and sometimes serology. Recent advances in technology make detection and identification faster, more convenient, more sensitive, and more specific than conventional assays -- at least in theory. These new methods are often referred to as "rapid methods", a subjective term used loosely to describe a vast array of tests that includes miniaturized biochemical kits, antibody- and DNA-based tests, and assays that are modifications of conventional tests to speed up analysis (8, 15, 16, 24, 36). Some of these assays have also been automated to reduce hands-on manipulations. With few exceptions, almost all assays used to detect specific pathogens in foods require some growth in an enrichment medium before analysis.

Experts who were surveyed in 1981 (19) about future developments in methods used for food microbiology, accurately predicted the widespread use of miniaturized biochemical kits for the identification of pure cultures of bacteria isolated from food. Most consist of a disposable device containing 15 - 30 media or substrates specifically designed to identify a bacterial group or species. With the exception of a few kits where results can be read in 4 hrs, most require 18-24 hrs incubation. In general, miniaturized biochemical tests are very similar in format and performance, showing 90-99% accuracy in comparison to conventional methods (5, 16, 21). However, kits that have been in use longer may have a more extensive identification database than newer tests. Most miniaturized kits are designed

for enteric bacteria, but kits for the identification of non-*Enterobacteriaceae* are also available, including for *Campylobacter*, *Listeria*, anaerobes, non-fermenting Gram-negative bacteria and for Gram-positive bacteria (Table 1).

Advances in instrumentation have enabled automation of the miniaturized biochemical identification tests. These instruments can incubate the reactions and automatically monitor biochemical changes to generate a phenotypic profile, which is then compared with the provided database stored in the computer to provide an identification (8, 23, 35). Other automated systems identify bacteria based on compositional or metabolic properties, such as fatty acid profiles, carbon oxidation profiles (28) or other traits (Table 1).

Not forecast in that 1981 survey were the potential applications of immunological and genetic techniques in food microbiology (19). During the 1980s, major advances in basic research were transferred rapidly to applied areas, as "biotechnology" companies emerged and sought markets in the diagnostic field (11). DNA and antibody-based assays for numerous microbes or their toxins are now available commercially (12).

There are many DNA-based assay formats, but only probes, PCR and bacteriophage have been developed commercially for detecting foodborne pathogens. Probe assays generally target ribosomal RNA (rRNA), taking advantage of the fact that the higher copy number of bacterial rRNA provides a naturally amplified target and affords greater assay sensitivity (6, 14, 25, 37) (Table 2).

The basic principle of DNA hybridization is also being utilized in other technologies, such as the polymerase chain reaction (PCR) assay, where short fragments of DNA (probes) or primers are hybridized to a specific sequence or template, which is then enzymatically amplified by *Taq* polymerase using a thermocycler (2, 22). Theoretically, PCR can amplify a single copy of DNA by a million fold in less than 2 hrs; hence its potential to eliminate, or greatly reduce the need for cultural enrichment. However, the presence of inhibitors in foods and in many culture media can prevent primer binding and diminish amplification efficiency (26, 34), so that the extreme sensitivity achievable by PCR with pure cultures is often reduced when testing foods. Therefore, some cultural enrichment is still required prior to analysis (Table 2).

The highly specific interaction of phage with its bacterial host has also been used to develop assays for foodborne pathogens (38). One example is an assay for *Salmonella*, in which a specific bacteriophage was engineered to carry a detectable marker (ice nucleation gene). In the presence of *Salmonella*, the phage confers the marker to the host, which then expresses the phenotype to allow detection (Table 2).

The highly specific binding of antibody to antigen, especially monoclonal antibody, plus the simplicity and versatility of this reaction, has facilitated the design of a variety of antibody assays and formats, and they comprise the largest group of rapid methods being used in food testing (3, 10, 12, 33). There are 5 basic formats of antibody assays (12), the simplest of which is latex agglutination (LA), in which antibody-coated colored latex beads or colloidal gold particles are used for quick serological identification or typing of pure culture isolates of bacteria from foods (7, 12). A modification of LA, known as reverse passive latex agglutination (RPLA), tests for soluble antigens and is used mostly in testing for toxins in food extracts or for toxin production by pure cultures (12) (Table 3).

In the immunodiffusion test format, an enrichment sample is placed in a gel matrix with the antibody; if the specific antigen is present, a visible line of precipitation is formed (30).

The enzyme-linked immunosorbent assay (ELISA) is the most prevalent antibody assay format used for pathogen detection in foods (3, 33). Usually designed as a "sandwich" assay, an antibody bound to a solid matrix is used to capture the antigen from enrichment cultures and a second antibody conjugated to an enzyme is used for detection. The walls of wells in microtiter plates are the most commonly used solid support; but ELISAs have also been designed using dipsticks, paddles, membranes, pipet tips or other solid matrices (12) (Table 3).

Antibodies coupled to magnetic particles or beads are also used in immunomagnetic separation (IMS) technology to capture pathogens from pre-enrichment media (31). IMS is analogous to selective enrichment, but instead of using antibiotics or harsh reagents that can cause stress-injury, an antibody is used to capture the antigen, which is a much milder alternative. Captured antigens can be plated or further tested using other assays.

Immunoprecipitation or immunochromatography, still another antibody assay format, is based on the technology developed for home pregnancy tests. It is also a "sandwich" procedure but, instead of enzyme conjugates, the detection antibody is coupled to colored latex beads or to colloidal gold. Using only a 0.1 ml aliquot, the enrichment sample is wicked across a series of chambers to obtain results (9). These assays are extremely simple, require no washing or manipulation and are completed within 10 minutes after cultural enrichment (Table 3).

The last mentioned "category" of rapid methods includes a large variety of assays, ranging from specialized media to simple modifications of conventional assays, which result in saving labor, time, and materials. Some, for instance, use disposable cardboards containing dehydrated media, which eliminates the need for agar plates, constituting savings in storage, incubation and disposal procedures (4, 5). Others incorporate specialized chromogenic and fluorogenic substrates in media to rapidly detect enzymatic activity (13, 17, 20, 27, 29). There are also tests that measure bacterial adenosine triphosphate (ATP), which (although not identifying specific species), can be used to rapidly enumerate the presence of total bacteria (Table 4).

Applications and Limitations of Rapid Methods

Almost all rapid methods are designed to detect a single target, which makes them ideal for use in quality control programs to quickly screen large numbers of food samples for the presence of a particular pathogen or toxin. A positive result by a rapid method however, is only regarded as presumptive and must be confirmed by standard methods (11). Although confirmation may extend analysis by several days, this may not be an imposing limitation, as negative results are most often encountered in food analysis.

Most rapid methods can be done in a few minutes to a few hours, so they are more rapid than traditional methods. But, in food analysis, rapid methods still lack sufficient sensitivity and specificity for direct testing; hence, foods still need to be culture-enriched before analysis (12). Although enrichment is a limitation in terms of assay speed, it provides essential benefits, such as diluting the effects of inhibitors, allowing the differentiation of viable from non-viable cells and allowing for repair of cell stress or injury that may have resulted during food processing.

Evaluations of rapid methods show that some perform better in some foods than others. This can be attributed mostly to interference by food components, some of which can be especially troublesome for the technologies used in rapid methods. For example, an ingredient can inhibit DNA hybridization or *Taq* polymerase, but has no effect on antigen-antibody interactions and the converse situation may also occur (12). Since method efficiencies may be food dependent, it is advisable to perform comparative studies to ensure that a particular assay will be effective in the analysis of that food type.

The specificity of DNA based assays is dictated by short probes; hence, a positive result, for instance with a probe or primers specific for a toxin gene, only indicates that bacteria with those gene sequences are present and that they have the potential to be toxigenic. But, it does not indicate that the gene is actually expressed and that the toxin is made. Likewise, in clostridial and staphylococcal intoxication, DNA probes and PCR can detect only the presence of cells, but are of limited use in detecting the presence of preformed toxins (12).

Currently, there are at least 30 assays each for testing for *E. coli* O157:H7 and for *Salmonella*. Such a large number of options can be confusing and overwhelming to the user, but, more importantly, has limited the effective evaluation of these methods. As a result, only few methods have been officially validated for use in food testing (1,11).

Conclusions

As a rapid method is used more frequently, its benefits and at the same time, its limitations also become more apparent. This section only briefly described some of the rapid method formats and selected problems encountered when using these assays in food analysis. However, because of the complex designs and formats of these tests, coupled with the difficulties of testing foods, users must exercise caution when selecting rapid methods and to also evaluate these tests thoroughly, as some may be more suitable than others for distinct testing situations or for assaying certain types of food. Lastly, technology continues to advance at a great pace and next generation assays, such as biosensors (18) and DNA chips (32) already are being developed that potentially have the capability for near real-time and on-line monitoring of multiple pathogens in foods.

NOTE: The listings provided in Tables 1-4 are intended for general reference only and do not indicate endorsement or approval by FDA for use in food analysis.

Table 1. Partial list of miniaturized biochemical kits and automated systems for identifying foodborne bacteria* (5, 8, 15, 16, 21, 35, 36).

System	Format	Manufacturer	Organisms
API ^b	biochemical	bioMerieux	<i>Enterobacteriaceae, Listeria, Staphylococcus, Campylobacter, Non-fermenters, anaerobes</i>
Cobas IDA	biochemical	Hoffmann LaRoche	<i>Enterobacteriaceae</i>
Micro-ID ^b	biochemical	REMEL	<i>Enterobacteriaceae, Listeria</i>
Enterotubell	biochemical	Roche	<i>Enterobacteriaceae</i>
Spectrum 10	biochemical	Austin Biological	<i>Enterobacteriaceae</i>
RapID	biochemical	Innovative Diag.	<i>Enterobacteriaceae</i>
BBL Crystal	biochemical	Becton Dickinson	<i>Enterobacteriaceae, Vibrionaceae, Non-fermenters, anaerobes</i>
Minitek	biochemical	Becton Dickinson	<i>Enterobacteriaceae</i>
Microbact	biochemical	Microgen	<i>Enterobacteriaceae, Gram negatives, Non-fermenters, Listeria</i>
Vitek ^b	biochemical ^a	bioMerieux	<i>Enterobacteriaceae, Gram negatives, Gram positives</i>
Microlog	C oxidation ^a	Biolog	<i>Enterobacteriaceae, Gram negatives, Gram positives</i>
MIS ^b	Fatty acid ^a	Microbial-ID	<i>Enterobacteriaceae, Listeria, Bacillus, Staphylococcus, Campylobacter</i>
Walk/Away	biochemical ^a	MicroScan	<i>Enterobacteriaceae, Listeria, Bacillus, Staphylococcus, Campylobacter</i>
Replianalyzer	biochemical ^a	Oxoid	<i>Enterobacteriaceae, Listeria, Bacillus, Staphylococcus, Campylobacter</i>
Riboprinter	nucleic acid ^a	Qualicon	<i>Salmonella, Staphylococcus, Listeria, Escherichia coli</i>
Cobas Micro-ID	biochemical ^a	Becton Dickinson	<i>Enterobacteriaceae, Gram negatives, Non-fermenters</i>
Malthus ^b	conductance ^a	Malthus	<i>Salmonella, Listeria, Campylobacter, E. coli, Pseudomonas, coliforms</i>
Bactometer	impedance ^a	bioMerieux	<i>Salmonella</i>

* Table modified from: Feng, P., App.I., FDA Bacteriological Analytical Manual, 8A ed.

^a Automated systems

^b Selected systems adopted AOAC Official First or Final Action.

NOTE: This table is intended for general reference only and lists known available methods. Presence on this list does not indicate verification, endorsement, or approval by FDA for use in food analysis.

Table 2. Partial list of commercially-available, nucleic acid-based assays used in the detection of foodborne bacterial pathogens* (2, 5, 8, 12, 14, 22, 25, 36, 37).

Organism	Trade Name	Format	Manufacturer
<i>Clostridium botulinum</i>	Probelia	PCR	BioControl
<i>Campylobacter</i>	AccuProbe	probe	GEN-PROBE
	GENE-TRAK	probe	Neogen
<i>Escherichia coli</i>	GENE-TRAK	probe	Neogen
<i>E. coli</i> O157:H7	BAX	PCR ^a	Qualicon
	Probelia	PCR	BioControl
<i>Listeria</i>	GENE-TRAK ^c	probe	Neogen
	AccuProbe	probe	GEN-PROBE
	BAX	PCR	Qualicon
	Probelia	PCR	BioControl
<i>Salmonella</i>	GENE-TRAK ^c	probe	Neogen
	BAX	PCR	Qualicon
	BIND ^b	phage	BioControl
	Probelia	PCR	BioControl
<i>Staphylococcus aureus</i>	AccuProbe	probe	GEN-PROBE
	GENE-TRAK	probe	Neogen
<i>Yersinia enterocolitica</i>	GENE-TRAK	probe	Neogen

* Table modified from: Feng, P., App.I, FDA Bacteriological Analytical Manual, 8A ed.
^a Polymerase chain reaction
^b Bacterial Ice Nucleation Diagnostics
^c Adopted AOAC Official First or Final Action

NOTE: This table is intended for general reference only and lists known available methods. Presence on this list does not indicate verification, endorsement, or approval by FDA for use in food analysis.

Table 3. Partial list of commercially-available, antibody-based assays for the detection of foodborne pathogens and toxins* (3, 5, 8, 12, 33, 36).

Organism/toxin	Trade Name	Assay Format ^a	Manufacturer
<i>Bacillus cereus</i> diarrhoeal toxin	TECRA	ELISA	TECRA
	BCET	RPLA	Unipath
<i>Campylobacter</i>	Campyslide	LA	Becton Dickinson
	Meritec-campy	LA	Meridian
	MicroScreen	LA	Mercia
	VIDAS	ELFA ^b	bioMerieux
	EiaFOSS	ELISA ^b	Foss
	TECRA	ELISA	TECRA
<i>Clostridium botulinum</i> toxin	ELCA	ELISA	Elcatech
<i>C. perfringens</i> enterotoxin	PET	RPLA	Unipath
<i>Escherichia coli</i>			
EHEC ^{**c} O157:H7	RIM	LA	REMEL
	<i>E. coli</i> O157	LA	Unipath
	Prolex	LA	PRO-LAB
	Ecolex O157	LA	Orion Diagnostica
	Wellcolex O157	LA	Murex
	<i>E. coli</i> O157	LA	TechLab
	O157&H7	sera	Difco
	PetrifilmHEC	Ab-blot	3M
	EZ COLI	Tube-EIA	Difco
	Dynabeads	Ab-beads	Dynal
	EHEC-TEK	ELISA	Organon-Teknika
	Assurance ^e	ELISA	BioControl
	HECO157	ELISA	3M Canada
	TECRA	ELISA	TECRA
	<i>E. coli</i> O157	ELISA	LMD Lab
	Premier O157	ELISA	Meridian
	<i>E. coli</i> O157:H7	ELISA	Binax
	<i>E. coli</i> Rapitest	ELISA	Microgen
	Transia Card <i>E. coli</i> O157	ELISA	Diffchamb
	<i>E. coli</i> O157	EIA/capture	TECRA
	VIP ^e	Ab-ppt	BioControl
Reveal	Ab-ppt	Neogen	

	Quix Rapid O157	Ab-ppt	Universal HealthWatch
	ImmunoCardSTAT	Ab-ppt	Meridian
	VIDAS	ELFA ^b	bioMerieux
	EiaFOSS	ELISA ^b	Foss
Shiga toxin (Stx)	VEROTEST	ELISA	MicroCarb
	Premier EHEC	ELISA	Meridian
	Verotox-F	RPLA	Denka Seiken
ETEC ^c			
Labile toxin (LT)	VET-RPLA	RPLA	Oxoid
Stabile toxin (ST)	E. coli ST	ELISA	Oxoid
<i>Listeria</i>	Microscreen	LA	Microgen
	Listeria Latex	LA	Microgen
	Listeria-TEK ^e	ELISA	Organon Teknika
	TECRA ^e	ELISA	TECRA
	Assurance ^e	ELISA	BioControl
	Transia Plate Listeria	ELISA	Diffchamb
	Pathalert	ELISA	Merck
	Listertest	Ab-beads	VICAM
	Dynabeads	Ab-beads	Dynal
	VIP ^e	Ab-ppt	BioControl
	Clearview	Ab-ppt	Unipath
	RAPIDTEST	Ab-ppt	Unipath
	VIDAS ^e	ELFA ^b	bioMerieux
	EiaFOSS	ELISA ^b	Foss
	UNIQUE	Capture-EIA	TECRA
<i>Salmonella</i>	Bactigen	LA	Wampole Labs
	Spectate	LA	Rhone-Poulenc
	Microscreen	LA	Mercia
	Wellcolex	LA	Laboratoire Wellcome
	Serobact	LA	REMEL
	RAPIDTEST	LA	Unipath
	Dynabeads	Ab-beads	Dynal
	Screen	Ab-beads	VICAM
	CHECKPOINT	Ab-blot	KPL
	1-2 Test ^e	diffusion	BioControl
	SalmonellaTEK ^e	ELISA	Organon Teknika
	TECRA ^e	ELISA	TECRA

	EQUATE	ELISA	Binax
	BacTrace	ELISA	KPL
	LOCATE	ELISA	Rhone-Poulenc
	Assurance ^e	ELISA	BioControl
	Salmonella	ELISA	GEM Biomedical
	Transia Plate Salmonella Gold	ELISA	Diffchamb
	Bioline	ELISA	Bioline
	VIDAS ^e	ELFA ^b	bioMerieux
	OPUS	ELISA ^b	TECRA
	PATH-STIK	Ab-ppt	LUMAC
	Reveal	Ab-ppt	Neogen
	Clearview	Ab-ppt	Unipath
	UNIQUE ^e	Capture-EIA	TECRA
<i>Shigella</i>	Bactigen	LA	Wampole Labs
	Wellcolex		Laboratoire Wellcome
<i>Staphylococcus aureus</i>	Staphyloslide	LA	Becton Dickinson
	AureusTest ^e	LA	Trisum
	Staph Latex	LA	Difco
	<i>S. aureus</i> VIA	ELISA	TECRA
enterotoxin	SET-EIA	ELISA	Toxin Technology
	SET-RPLA	RPLA	Unipath
	TECRA ^e	ELISA	TECRA
	Transia Plate SE	ELISA	Diffchamb
	RIDASCREEN	ELISA	R-Biopharm
	VIDAS	ELFA ^b	bioMerieux
	OPUS	ELISA ^b	TECRA
<i>Vibrio cholera</i>	choleraSMART	Ab-ppt	New Horizon
	bengalSMART	Ab-ppt	New Horizon
	choleraScreen	Agglutination	New Horizon
	bengalScreen	Agglutination	New Horizon
enterotoxin	VET-RPLA ^d	RPLA	Unipath
<p>* Table modified from: Feng, P., App.I, FDA Bacteriological Analytical Manual, 8A ed. ^a Abbreviations: ELISA, enzyme linked immunosorbent assay; ELFA, enzyme linked fluorescent assay; RPLA, reverse passive latex agglutination; LA, latex agglutination; Ab-ppt, immunoprecipitation. ^b Automated ELISA ^c EHEC - Enterohemorrhagic <i>E. coli</i>; ETEC - enterotoxigenic <i>E. coli</i> ^d Also detects <i>E. coli</i> LT enterotoxin ^e Adopted AOAC Official First or Final Action</p>			
** CAUTION: unless the assays claim that they are specific for the O157:H7 serotype, most of			

these tests detect only the O157 antigen; hence will also react with O157 strains that are not of H7 serotype. These O157, non-H7 strains, generally do not produce Shiga toxins and are regarded as not pathogenic for humans. Furthermore, some antibodies to O157 can also cross react with *Citrobacter*, *E. hermanii* and other enteric organisms.

NOTE: This table is intended for general reference only and lists known available methods. Presence on this list does not indicate verification, endorsement, or approval by FDA for use in food analysis.

Table 4. Partial list of other commercially available rapid methods and specialty substrate media for detection of foodborne bacteria* (4, 8, 13, 20, 27, 36).

Organism	Trade Name	Format ^a	Assay Manufacturer
Bacteria	Redigel ^b	Media	RCR Scientific
	Isogrid ^b	HGMF	QA Labs
	Enliten	ATP	Promega
	Profile-1	ATP	New Horizon
	Biotrace	ATP	Biotrace
	Lightning	ATP	Idexx
	Petrifilm ^b	media-film	3M
	Sim Plate	media	Idexx
Coliform/ <i>E. coli</i>	Isogrid ^b	HGMF/MUG	QA Labs
	Petrifilm ^b	media-film	3M
	SimPlate	media	Idexx
	Redigel	Media	RCR Scientific
	ColiQuik ^c	MUG/ONPG	Hach
	ColiBlue ^c	media	Hach
	Colilert ^{b,c}	MUG/ONPG	Idexx
	LST-MUG ^b	MPN media	Difco & GIBCO
	ColiComplete ^b	MUG-Xgal	BioControl
	Colitrak	MPN-MUG	BioControl
	ColiGel & E*Colite ^c	MUG-Xgal	Charm Sciences
	CHROMagar	Medium	CHROMagar
	<i>E. coli</i>	MUG disc	MUG
CHROMagar		Medium	CHROMagar
EHEC ^d	Rainbow Agar	Medium	Biolog
	BCMO157:H7	Medium	Biosynth
	Fluorocult O157:H7	Medium	Merck
<i>Listeria monocytogenes</i>	BCM	Medium	Biosynth
<i>Salmonella</i>	Isogrid ^b	HGMF	QA Labs
	OSRT	Medium/ motility	Unipath (Oxoid)
	Rambach	Medium	CHROMagar
	MUCAP	C8esterase	Biolife
	XLT-4	Medium	Difco
	MSRV ^b	Medium	
<i>Yersinia</i>	Crystal violet	Dye binding	Polysciences

* Table modified from: Feng, P., App.I, FDA Bacteriological Analytical Manual, 8A ed.

^a Abbreviations: APC, aerobic plate count; HGMF, hydrophobic grid membrane filter; ATP, adenosine triphosphate; MUG, 4-methylumbelliferyl- β -D-glucuronide; ONPG, O-nitrophenyl β -D-galactoside; MPN, most probable number.

^b Adopted AOAC Official First or Final Action.

^c Application for water analysis

^d EHEC - enterohemorrhagic *Escherichia coli*

NOTE: This table is intended for general reference only and lists known available methods. Presence on this list does not indicate verification, endorsement, or approval by FDA for use in food analysis.

References

1. Andrews, W.H. 1996. AOAC INTERNATIONAL'S three validation programs for methods used in the microbiological analysis of foods. *Trend in Food Sci. Technol.* **7**:147-151.
2. Barrett, T., P. Feng and B. Swaminathan. 1997. Amplification Methods for Detection of Foodborne Pathogens, pp.171-181. *In*: H.H. Lee, S.A. Morse and O. Olsvik (ed), *Nucleic Acid Amplification Techniques: Application to Disease Diagnosis*. Eaton Publishing, Boston.
3. Candish, A.A.G. 1991. Immunological methods in food microbiology. *Food Microbiol.* **8**:1-14.
4. Chain, V.S., and D.Y.C. Fung. 1991. Comparison of Redigel, Petrifilm, spiral plate system, Isogrid, and aerobic plate count for determining the numbers of aerobic bacteria in selected foods. *J. Food Prot.* **54**:208-211.
5. Cox, N.A., D.Y.C. Fung, J.S. Bailey, P.A. Hartman, and P.C. Vasavada. 1987. Miniaturized kits, immunoassays and DNA hybridization for recognition and identification of foodborne bacteria. *Dairy Food Sanit.* **7**:628-631.
6. Curiale, M.S., M.J. Klatt, and M.A. Mozola. 1990. Colorimetric deoxyribonucleic acid hybridization assay for rapid screening of *Salmonella* in foods: Collaborative study. *J. Assoc. Off. Anal. Chem.* **73**:248-256.
7. D'Aoust, J.-Y., A.M. Sewell, and P. Greco. 1991. Commercial latex agglutination kits for the detection of foodborne *Salmonella*. *J. Food Prot.* **54**:725-730.
8. Dziezak, J.D. 1987. Rapid methods for microbiological analysis of foods. *Food Technol.* **41**(7):56-73.
9. Feldsine, P.T., R.L. Forgey, M.T. Falbo-Nelson and S. Brunelle. 1997. *Escherichia coli* O157:H7 Visual Immunoprecipitation assay: a comparative validation study. *J. AOAC* **80**:43-48.
10. Feng, P. 1992. Commercial assay systems for detecting foodborne *Salmonella*: a review. *J. Food Prot.* **55**:927-934.
11. Feng, P. 1996. Emergence of rapid methods for identifying microbial pathogens in foods. *J. Assoc. Off. Anal. Chem. Int.* **79**:809-812.
12. Feng, P. 1997. Impact of Molecular Biology on the Detection of Foodborne Pathogens. *Mol. Biotech.* **7**:267-278.
13. Feng, P.C.S., and P.A. Hartman. 1982. Fluorogenic assays for immediate confirmation of *Escherichia coli*. *Appl. Environ. Microbiol.* **43**:1320-1329.
14. Feng, P., K.A. Lampel and W.E. Hill. 1996. Developments in food technology: applications and economic and regulatory considerations, pp. 203-229. *In*: C.A. Dangler and B. Osburn (ed). *Nucleic Acid Analysis: Principles and Bioapplications*. Wiley & Sons, New York.
15. Fung, D.Y.C. 1991. Rapid methods and automation for food microbiology, pp. 1-38. *In*: *Instrumental Methods for Quality Assurance in Foods*. D.Y.C. Fung and R.F. Matthews (eds). Marcel Dekker, New York.
16. Fung, D.Y.C., N.A. Cox, and J.S. Bailey. 1988. Rapid methods and automation in the microbiological examination of food. *Dairy Food Sanit.* **8**:292-296.
17. Gaillot, O., P.D. Camillo, P. Berche, R. Courcol and C. Savage. 1999. Comparison of CHROMagar salmonella medium and Hektoen Enteric agar for isolation of salmonellae from stool samples. *J. Clin. Microbiol.* **37**:762-765.
18. Goldschmidt, M. 1999. Biosensors - scope in microbiological analysis, p. 268-278. *In* R.K. Robinson, C.A. Batt and P. Patel (ed.). *Encyclopedia of Food Microbiology*. Academic Press, London.
19. Gutteridge, C.S., and M.L. Arnott. 1989. Rapid methods: an over the horizon view, pp. 297-319. *In*: *Rapid Methods in Food Microbiology: Progress in Industrial Microbiology*. M.R. Adams and C.F.A. Hope (eds). Elsevier, New York.
20. Hartman, P.A. 1989. The MUG (glucuronidase) test for *Escherichia coli* in food and water, pp. 290-308. *In*: *Rapid Methods and Automation in Microbiology and Immunology*. A. Balows, R.C. Tilton, and A. Turano (eds). Brixia Academic Press, Brescia, Italy.
21. Hartman, P.A., B. Swaminathan, M.S. Curiale, R. Firstenberg-Eden, A.N. Sharpe, N.A. Cox, D.Y.C. Fung, and M.C. Goldschmidt. 1992. Chapter 39, Rapid methods and automation, pp. 665-746. *In*: *Compendium of Methods for the Microbiological Examination of Foods*, 3rd ed., C. Vanderzant and D.F. Splittstoesser (eds). American Public Health Association, Washington, DC.
22. Hill, W.E. 1996. The polymerase chain reaction: application for the detection of foodborne pathogens. *CRC Crit. Rev. Food Sci. Nutr.* **36**:123-173.

23. Holmes, B. 1989. Comparative evaluation of the Roche Cobas IDA and Enterotube II systems for identifying members of the family *Enterobacteriaceae*. *J. Clin. Microbiol.* 27:1027-1030.
24. Ibrahim, G.F. 1986. A review of immunoassays and their application to salmonellae detection in foods. *J. Food Prot.* 49:299-310.
25. Kalamaki, M., R.J. Pricce and D.Y.C. Fung. 1997. Rapid methods for identifying seafood microbial pathogens and toxins. *J. Rapid Methods and Automation in Microbiol.* 5:87-137.
26. Lampel, K.A., P. Feng and W.E. Hill. 1992. Gene probes used in food microbiology, pp. 151-188. *In* D. Bhatnagar and T.E. Cleveland (ed.), *Molecular Approaches to Improving Food Safety*. Van Nostrand Reinhold, New York, NY.
27. Lantz, P-G., B. Hahn-Hagerdal, and P. Radstrom. 1994. Sample preparation methods in PCR-based detection of food pathogens. *Trend Food Sci. Technol.* 5:384-389.
28. Manafi, M., W. Kneifel and S. Bascomb. 1991. Fluorogenic and chromogenic substrates used in bacterial diagnostics. *Microbiol. Rev.* 55:335-348.
29. Miller, J.M., and D.L. Rhoden. 1991. Preliminary evaluation of Biolog, a carbon source utilization method for bacterial identification. *J. Clin. Microbiol.* 29:1143-1147.
30. Moberg, L.J., M.K. Wagner, and L.A. Kellen. 1988. Fluorogenic assay for rapid detection of *Escherichia coli* in chilled and frozen foods: collaborative study. *J. Assoc. Off. Anal. Chem.* 71:589-602.
31. Oggel, J.J., D.C. Nundy, and C.J. Randall. 1990. Modified 1-2 test system as a rapid screening method for detection of *Salmonella* in foods and feeds. *J. Food Prot.* 53:656-658.
32. Olsvik, O., T. Popovic, E. Skjerve, K.S. Cudjoe, E. Hornes, J. Ugelstad and M. Uhlen. 1994. Magnetic separation techniques in diagnostic microbiology. *Clin. Microbiol. Rev.* 7:43-54.
33. Ramsay, G. 1998. DNA chips: State-of-the art. *Nature Biotechnol.* 16:40-44.
34. Rose, S.A., and M.F. Stringer. 1989. Immunological methods, pp. 121-167. *In*: *Rapid Methods in Food Microbiology: Progress in Industrial Microbiology*. M.R. Adams and C.F.A. Hope (eds). Elsevier, New York.
35. Rossen, L., P. Norskov, K. Holmstrom, O.F. Rasmussen. 1992. Inhibition of PCR by components of food samples, and DNA-extraction solutions. *Int. J. Food Microbiol.* 17:37-45.
36. Stager, C.E., and J.R. Davis. 1992. Automated systems for identification of microorganisms. *Clin. Microbiol. Rev.* 5:302-327.
37. Swaminathan, B. and P. Feng. 1994. Rapid Detection of Foodborne Pathogenic Bacteria. *In*: *Annual Review of Microbiology*, 48:401-426. Annual Reviews Inc., Palo Alto, CA.
38. Tenover, F.C. 1988. Diagnostic deoxyribonucleic acid probes for infectious diseases. *Clin. Microbiol. Rev.* 1:82-101.
39. Wolber, P.K. and R.L. Green. 1990. New rapid method for the detection of *Salmonella* in foods. *Trends Food Sci. Technol.* 1:80-82.

Hypertext Source: Bacteriological Analytical Manual, 8th Edition, Revision A, 1998. Revised 2000-July-18, Final Revision: 2001-Jan-25

Author: [Peter Feng](#)