

HFD 104

Murphy

APR 28 1999

NDA 19-839

Pfizer Pharmaceuticals  
Attention: Martha Brumfield, Ph.D.  
Drug Regulatory Affairs  
235 East 42nd Street  
New York, New York 10017-3184

Three Years from APR 28 2002  
The Date of This Letter

Dear Dr. Brumfield:

Reference is made to your Proposed Pediatric Study Request submitted on August 31, 1998 to your Investigational New Drug (IND) application for Zoloft (sertraline hydrochloride) tablets

(b) (4)

We have completed our review of your submission and concluded that your proposed pediatric study request is incomplete.

To obtain needed pediatric information on sertraline, the Food and Drug Administration (FDA) is hereby making a formal Written Request, pursuant to Section 505A of the Federal Food, Drug, and Cosmetic Act (the Act), that you submit information from the trials in pediatric patients with depression described below.

(b) (4)

**Specific Study Requirements**

(b) (4)

**Types of Studies**

Pediatric Efficacy and Safety Studies

Pediatric Pharmacokinetic Study

Pediatric Safety Study

**Objective/Rationale**

The overall goal (b) (4) is to establish the safety and efficacy of the study drug in the treatment of pediatric depression, and to develop other information, e.g., pharmacokinetic, pertinent to using the drug in the pediatric population.

**Study Design**

Pediatric Efficacy and Safety Studies

- For the controlled efficacy studies, conduct two randomized, double-blind, parallel group, placebo-controlled acute treatment trial, with a recommended duration of at least 6 to 8 weeks. We recommend that at least one of the two studies should be a fixed dose study including two or more fixed doses of the study drug. You may consider dosing patients on the basis of patient weight. Randomization must be stratified by the two age groups studied. Ideally, a relapse prevention trial would follow from the acute treatment trials, involving the randomization of responders from the acute treatment trials to continuation on either study drug or placebo, with follow-up observation for relapse for a period of 6 months or more. Please note that a relapse prevention trial is not required under this written request.

Pediatric Pharmacokinetic Study

- A pharmacokinetic study to provide information pertinent to dosing of the study drug in the relevant pediatric population. These data could come from traditional pharmacokinetic studies, or alternatively, from population kinetic approaches applied to controlled efficacy trials or to other safety trials. You should be aware that a guidance document on population pharmacokinetic studies is available under [[www.fda.gov/cder/guidance/1852fml.pdf](http://www.fda.gov/cder/guidance/1852fml.pdf)].

Pediatric Safety Study

- Safety data should be collected in the controlled efficacy trials. Longer-term safety data should be generated in longer-term open extensions from these trials and/or in separate longer-term open safety studies.

**Age Group in Which Study(ies) will be Performed – All Studies**

Both children (ages 7 to 11) and adolescents (ages 12 to 17) should be equally represented in the samples, and there should be a reasonable distribution of both sexes in these strata.

**Number of Patients to be Studied or Power of Study to be Achieved****Pediatric Efficacy and Safety Studies**

- While it is difficult to specify the sample size needed to show a difference between drug and placebo in this population, it should be noted that, in the only published positive antidepressant trial in pediatric depression (Emslie, et al, 1997), there were 48 patients in each of the two treatment arms.

**Pediatric Pharmacokinetic Study**

- A sufficient number of subjects to adequately characterize the pharmacokinetics in the above age groups.

**Pediatric Safety Study**

- A sufficient number of pediatric patients to adequately characterize the safety of sertraline at clinically effective doses for a sufficient duration.

**Entry Criteria**

The protocols should include a valid and reliable diagnostic method for recruiting children and adolescents with major depressive disorder.

**Study Endpoints****Pediatric Efficacy and Safety Studies**

- It is essential to identify a single primary outcome for the controlled efficacy trials, and ordinarily this should be change from baseline to endpoint on whatever symptom rating scale you have chosen for your trials.

**Pediatric Pharmacokinetic Study**

- Pharmacokinetic measurements as appropriate.

**Pediatric Safety Study**

- Appropriately frequent standard measures of safety (clinical - including signs and symptoms and laboratory).

**Statistical Information****Pediatric Efficacy and Safety Studies**

- These trials should have a detailed statistical plan. Ordinarily these trials should be designed with at least 80% statistical power to detect a treatment effect of conventional ( $p=0.05$ ) statistical significance.

**Pediatric Pharmacokinetic Study**

- Descriptive analysis of the pharmacokinetic parameters.

**Pediatric Safety Study**

- Descriptive analysis of the safety data.

**Study Evaluations****Pediatric Efficacy and Safety Studies**

- A scale specific to pediatric depression and sensitive to the effects of drug treatment of pediatric depression, e.g., the Children's Depression Rating Scale—Revised, and a global measure, e.g., the Clinical Global Impression (CGI).

**Pediatric Pharmacokinetic Study**

- The pharmacokinetic assessments should be made with respect to the study drug and any metabolites that make substantial contributions to its efficacy and/or toxicity. For the parent and each metabolite followed, the data collected should provide estimates of the pharmacokinetic parameters including AUC, half-life,  $C_{max}$ ,  $t_{max}$ , and apparent oral clearance in pediatric subjects in the relevant age range. You should be aware that a draft guidance document on pediatric pharmacokinetic studies is available under [[www.fda.gov/cder/guidance/index.htm](http://www.fda.gov/cder/guidance/index.htm), under Clinical/Pharmacological (Draft)].

**Pediatric Safety Study**

- Routine safety assessments should include vital signs, weight, clinical laboratory, ECGs, and monitoring for adverse events. Although not a part of this Written Request, we remind you that it may be important to determine the effect of the study drug on the growth and development of pediatric patients, and we encourage you to consider longer-term studies of a year or more to address this question if the acute studies demonstrate antidepressant activity.

**Drug Information**

Use age appropriate formulations in the studies described above. Since the pediatric patient population consists of both children (ages 7 to 11) and adolescents (ages 12 to 17), your marketed solid dosage formulation should be adequate for these studies.

**Drug Concerns**

No specific concerns related to administration to pediatric patients were identified while studying sertraline in adults, nor have specific concerns been identified during the postmarketing experience.

**Labeling That May Result from the Studies**

Results found in the pediatric depression population efficacy, safety, and pharmacokinetic studies could result in the addition to labeling of information pertinent to these studies.

**Format of Reports to be Submitted**

Full study reports or analyses, not previously submitted to the Agency, addressing the issues outlined in this request, with full analysis, assessment, and interpretation.

**Timeframe for Submitting Reports of the Study(ies)**

Reports of the above studies must be submitted to the Agency within 3 years from the date of this letter to be eligible to qualify for pediatric exclusivity extension under Section 505A of the Act. Please remember that pediatric exclusivity extends only existing patent protection or exclusivity that has not expired at the time you submit your reports of studies in response to this Written Request.

Please submit protocols for the above studies to an investigational new drug application (IND) and clearly mark your submission "**PEDIATRIC PROTOCOL SUBMITTED FOR PEDIATRIC EXCLUSIVITY STUDY**" in large font, bolded type at the beginning of the cover letter of the submission. We recommend you seek a written agreement with FDA before developing pediatric studies. Please notify us as soon as possible if you wish to enter into a written agreement by submitting a proposed written agreement. Clearly mark your submission "**PROPOSED WRITTEN AGREEMENT FOR PEDIATRIC STUDIES**" in large font, bolded type at the beginning of the cover letter of the submission.

Reports of the studies should be submitted as a supplement to your approved NDA with the proposed labeling changes you believe would be warranted based on the data derived from these studies. When submitting the reports, please clearly mark your submission "**SUBMISSION OF PEDIATRIC STUDY REPORTS – PEDIATRIC EXCLUSIVITY DETERMINATION REQUESTED**" in large font, bolded type at the beginning of the cover letter of the submission and include a copy of this letter. Please also send a copy of the cover letter of your submission, via fax (301-594-0183) or messenger to the Director, Office of Generic Drugs, HFD-600, Metro Park North II, 7500 Standish Place, Rockville, MD 20855-2773.

If you wish to discuss any amendments to this Written Request, please submit proposed changes and the reasons for the proposed changes to your application. Submissions of proposed changes to this request should be clearly marked "**PROPOSED CHANGES IN WRITTEN REQUEST FOR PEDIATRIC STUDIES**" in large font, bolded type at the beginning of the cover letter of the submission. You will be notified in writing if any changes to this Written Request are agreed upon by the Agency.

We hope you will fulfill this pediatric study request. We look forward to working with you on this matter in order to develop additional pediatric information that may produce health benefits to the pediatric population.

If you have any questions, contact Paul A. David, Regulatory Project Manager, at (301) 594-5530.

Sincerely yours,

A handwritten signature in black ink that reads "Robert Temple" followed by a date "4/28/98". The signature is written in a cursive, flowing style.

Robert Temple, M.D.  
Director  
Office of Drug Evaluation I  
Center for Drug Evaluation and Research

cc:

Archival NDA 19-839

HFD-120/division file

HFD-120/PDavid

HFD-120/RKatz/TLaughren/Amosholder/RGlass

HFD-100/RTemple

HFD-600/Office of Generic Drugs

HFD-2/MLumpkin

HFD-104/DMurphy

HFD-6/ KRoberts

Drafted:03/22/99am;rg

Rev:03/29/99tl; 04/02/99rt;04/12/99tl;4/21/99pdit

Final:04/22/99pd

Final: filename: DAVID\PEDIATRICS\ZOLOFTPEDS02.DOC

PEDIATRIC WRITTEN REQUEST LETTER

INFORMATION REQUEST (IR)